

Replica Bethe ansatz derivation of the Tracy-Widom distribution of the free energy fluctuations in one-dimensional directed polymers

Victor Dotsenko^{a,b}

^a*LPTMC, Université Paris VI, 75252 Paris, France and*

^b*L.D. Landau Institute for Theoretical Physics, 119334 Moscow, Russia*

(Dated: April 27, 2010)

The distribution function of the free energy fluctuations in one-dimensional directed polymers with δ -correlated random potential is studied by mapping the replicated problem to the N -particle quantum boson system with attractive interactions. We find the full set of eigenfunctions and eigenvalues of this many-body system and perform the summation over the entire spectrum of excited states. It is shown that in the thermodynamic limit the problem is reduced to the Fredholm determinant with the Airy kernel yielding the universal Tracy-Widom distribution, which is known to describe the statistical properties of the Gaussian unitary ensemble as well as many other statistical systems.

PACS numbers: 05.20.-y 75.10.Nr 74.25.Qt 61.41.+e

Contents

I. Introduction	 1
II. Mapping to quantum bosons	 3
III. Replica partition function and the free energy distribution function	 5
IV. Conclusions	 7
 Appendix A: Quantum bosons with repulsive interactions	
1. Eigenfunctions	 8
2. Othonormality	 9
Appendix B: Quantum bosons with attractive interactions	
1. Ground state	 10
2. Eigenfunctions	 12
3. Othonormality	 14
4. Propagator	 18
Appendix C: Fredholm determinant with the Airy kernel and the Tracy-Widom distribution	 20

I. INTRODUCTION

Directed polymers in a quenched random potential have been the subject of intense investigations during the past two decades (see e.g. [1]). Diverse physical systems such as domain walls in magnetic films [2], vortices in superconductors [3], wetting fronts on planar systems [4], or Burgers turbulence [5] can be mapped to this model, which exhibits numerous non-trivial features deriving from the interplay between elasticity and disorder. In the most simple one-dimensional case we deal with an elastic string directed along the τ -axis within an interval $[0, L]$. Randomness enters the problem through a disorder potential $V[\phi(\tau), \tau]$, which competes against the elastic energy. The problem is defined by the Hamiltonian

$$H[\phi(\tau), V] = \int_0^L d\tau \left\{ \frac{1}{2} [\partial_\tau \phi(\tau)]^2 + V[\phi(\tau), \tau] \right\}; \quad (1)$$

where in the simplest case the disorder potential $V[\phi, \tau]$ is Gaussian distributed with a zero mean $\overline{V(\phi, \tau)} = 0$ and the δ -correlations:

$$\overline{V(\phi, \tau)V(\phi', \tau')} = u\delta(\tau - \tau')\delta(\phi - \phi') \quad (2)$$

Here the parameter u describes the strength of the disorder. Historically, the problem of central interest was the scaling behavior of the polymer mean squared displacement which in the thermodynamic limit ($L \rightarrow \infty$) is believed to have a universal scaling form $\overline{\langle \phi^2 \rangle}(L) \propto L^{2\zeta}$ (where $\langle \dots \rangle$ and $\overline{(\dots)}$ denote the thermal and the disorder averages), with $\zeta = 2/3$, the so-called wandering exponent. More general problem for all directed polymer systems of the type, Eq.(1), is the statistical properties of their free energy fluctuations. Besides the usual extensive (linear in L) self-averaging part $f_0 L$ (where f_0 is the linear free energy density), the total free energy F of such systems contains disorder dependent fluctuating contribution $\tilde{F}$, which is characterized by non-trivial scaling in L . It is generally believed that in the limit of large L the typical value of the free energy fluctuations scales with L as $\tilde{F} \propto L^{1/3}$ (see e.g. [6–9]). In other words, in the limit of large L the total (random) free energy of the system can be represented as

$$F = f_0 L + c L^{1/3} f \quad (3)$$

where c is the parameter, which depends on the temperature and the strength of disorder, and f is the random quantity which in the thermodynamic limit $L \rightarrow \infty$ is described by a non-trivial universal distribution function $P_*(f)$. The derivation of this function for the system with δ -correlated random potential, Eqs.(1)-(2) is the central issue of the present work.

For the string with the zero boundary conditions at $\tau = 0$ and at $\tau = L$ the partition function of a given sample is

$$Z[V] = \int_{\phi(0)=0}^{\phi(L)=0} \mathcal{D}[\phi(\tau)] e^{-\beta H[\phi, V]} \quad (4)$$

where β denotes the inverse temperature. On the other hand, the partition function is related to the total free energy $F[V]$ via

$$Z[V] = \exp(-\beta F[V]) \quad (5)$$

The free energy $F[V]$ is defined for a specific realization of the random potential V and thus represent a random variable. Taking the N -th power of both sides of this relation and performing the averaging over the random potential V we obtain

$$\overline{Z^N[V]} \equiv Z[N, L] = \overline{\exp(-\beta N F[V])} \quad (6)$$

where the quantity in the lhs of the above equation is called the replica partition function. Substituting here $F = f_0 L + c L^{1/3} f$, and redefining

$$Z[N, L] = \tilde{Z}[N, L] e^{-\beta N f_0 L} \quad (7)$$

we get

$$\tilde{Z}[N, L] = \overline{\exp(-\lambda N f)} \quad (8)$$

where $\lambda = \beta c L^{1/3}$. The averaging in the rhs of the above equation can be represented in terms of the distribution function $P_L(f)$ (which depends on the system size L). In this way we arrive to the following general relation between the replica partition function $\tilde{Z}[N, L]$ and the distribution function of the free energy fluctuations $P_L(f)$:

$$\tilde{Z}[N, L] = \int_{-\infty}^{+\infty} df P_L(f) e^{-\lambda N f} \quad (9)$$

Of course, the most interesting object is the thermodynamic limit distribution function $P_*(f) = \lim_{L \rightarrow \infty} P_L(f)$ which is expected to be the universal quantity. The above equation is the bilateral Laplace transform of the function $P_L(f)$, and at least formally it allows to restore this function via inverse Laplace transform of the replica partition function $\tilde{Z}[N, L]$. In order to do so one has to compute $\tilde{Z}[N, L]$ for an arbitrary integer N and then perform analytical continuation of this function from integer to arbitrary complex values of N . In Kardar's original solution [9], after mapping the replicated problem to interacting quantum bosons, one arrives at the replica partition function for positive integer parameters $N > 1$. Assuming a large $L \rightarrow \infty$ limit, one is tempted to approximate the result by the ground state contribution only, as for any $N > 1$ the contributions of excited states are exponentially small for $L \rightarrow \infty$. However, in the analytic continuation for arbitrary complex N the contributions which are exponentially small at positive integer $N > 1$ can become essential in the region $N \rightarrow 0$, which defines the function $P(f)$ (in other word, the problem is that the two limits $L \rightarrow \infty$ and $N \rightarrow 0$ do not commute [10, 11]). As a consequence, the resulting distribution function $P_*(f)$ exhibits severe pathologies such as the vanishing of its second moment, which assumes that

the distribution function is not positively defined. Nevertheless, in terms of this approximation (assuming the universal scaling $L^{1/3}$ of the free energy fluctuations) one can derive the left tail of the distribution function $P_*(f)$ which is given by the asymptotics of the Airy function, $P_*(f \rightarrow -\infty) \sim \exp(-\frac{2}{3}|f|^{3/2})$ [12]. For the first time the form of the *right tail* of this distribution function, $P_*(f \rightarrow +\infty) \sim \exp[-(const)f^3]$, has been derived in terms of the optimal fluctuation approach [13], where it has been demonstrated that both asymptotics (left and right) of the function $P_*(f)$ are consistent with the Tracy-Widom (TW) distribution [14]. Originally the TW distribution has been derived in the context of the statistical properties of the Gaussian Unitary Ensemble while at present it is well established to describe the statistics of fluctuations in various random systems [15–20], which were widely believed to belong to the same universality class as the present model [13, 21, 22]. Finally, quite recently, the one-point distribution of the solutions of the KPZ-equation (which is equivalent to the present model) has been derived for an arbitrary value of L , and it has been shown that in the limit $L \rightarrow \infty$ this distribution turns into the Tracy-Widom distribution [23, 24].

In the recent paper [25] an attempt has been made to derive the free energy distribution function $P_*(f)$ via the calculation of the replica partition function $Z[N, L]$ in terms of the Bethe-Ansatz solution for quantum bosons with attractive δ -interactions which involved the summation over the *entire spectrum* of excited states. Unfortunately, the attempt has failed because on one hand, the calculations contained a kind of a hidden "uncontrolled approximation", and on the other hand, the analytic continuation of obtained $Z(N, L)$ was found to be ambiguous. Later on it turned out that it is possible to bypass the problem of the analytic continuation if instead of the distribution function itself one would study its integral representation

$$W(x) = \int_x^\infty df P_*(f) \quad (10)$$

which gives the probability to find the fluctuation f bigger than a given value x . Formally the function $W(x)$ can be defined as follows:

$$W(x) = \lim_{\lambda \rightarrow \infty} \sum_{N=0}^{\infty} \frac{(-1)^N}{N!} \exp(\lambda N x) \overline{\tilde{Z}^N} = \lim_{\lambda \rightarrow \infty} \overline{\exp[-\exp(\lambda(x-f))]} = \overline{\theta(f-x)} \quad (11)$$

On the other hand, the probability function $W(x)$ can be computed in terms of the replica partition function $\tilde{Z}[N, L]$ by summing over all replica integers

$$W(x) = \lim_{\lambda \rightarrow \infty} \sum_{N=0}^{\infty} \frac{(-1)^N}{N!} \exp(\lambda N x) \tilde{Z}[N, L] \quad (12)$$

In terms of this approach, quite recently it has been reported that performing the summation over the entire spectrum of excited states of N -particle quantum boson system the above series, eq.(12), can be reduced to the Fredholm determinant with the Airy kernel which is known to yield the Tracy-Widom distribution [26, 27].

Present paper provides the detailed derivation of this result. In particular, Appendices A and B contain detailed analysis of the properties of N -particle wave functions of quantum bosons with both repulsive and attractive δ -interactions. Besides, basing on the definition of the Fredholm determinant with the Airy kernel, in Appendix C it is demonstrated how the explicit form of the Tracy-Widom distribution can be derived in terms of the solution of the Painlevé II differential equation.

II. MAPPING TO QUANTUM BOSONS

Explicitly, the replica partition function, Eq.(6), of the system described by the Hamiltonian, Eq.(1), is

$$Z(N, L) = \prod_{a=1}^N \int_{\phi_a(0)=0}^{\phi_a(L)=0} \mathcal{D}\phi_a(\tau) \exp \left[-\beta \int_0^L d\tau \sum_{a=1}^N \left\{ \frac{1}{2} [\partial_\tau \phi_a(\tau)]^2 + V[\phi_a(\tau), \tau] \right\} \right] \quad (13)$$

Since the random potential $V[\phi, \tau]$ has the Gaussian distribution the disorder averaging ($\overline{\dots}$) in the above equation is very simple:

$$\overline{\exp \left[-\beta \int_0^L d\tau \sum_{a=1}^N V[\phi_a(\tau), \tau] \right]} = \exp \left[\frac{\beta^2}{2} \int_0^L \int_0^L d\tau d\tau' \sum_{a,b=1}^N \overline{V[\phi_a(\tau), \tau] V[\phi_b(\tau'), \tau']} \right] \quad (14)$$

Using Eq.(2) we find:

$$Z(N, L) = \prod_{a=1}^N \int_{\phi_a(0)=0}^{\phi_a(L)=0} \mathcal{D}\phi_a(\tau) \exp \left[-\frac{1}{2}\beta \int_0^L d\tau \left\{ \sum_{a=1}^N [\partial_\tau \phi_a(\tau)]^2 - \beta u \sum_{a,b=1}^N \delta[\phi_a(\tau) - \phi_b(\tau)] \right\} \right] \quad (15)$$

It should be noted that the second term in the exponential of the above equation contain formally divergent contributions proportional to $\delta(0)$ (due to the terms with $a = b$). In fact, this is just an indication that the *continuous* model, Eqs.(1)-(2) is ill defined as short distances and requires proper lattice regularization. Of course, the corresponding lattice model would contain no divergences, and the terms with $a = b$ in the exponential of the corresponding replica partition function would produce irrelevant constant $\frac{1}{2}L\beta^2 u N \delta(0)$ (where the lattice version of $\delta(0)$ has a finite value). Since the lattice regularization has no impact on the continuous long distance properties of the considered system this term will be just omitted in our further study.

Introducing the N -component scalar field replica Hamiltonian

$$H_N[\phi] = \frac{1}{2} \int_0^L d\tau \left(\sum_{a=1}^N [\partial_\tau \phi_a(\tau)]^2 - \beta u \sum_{a \neq b}^N \delta[\phi_a(\tau) - \phi_b(\tau)] \right) \quad (16)$$

for the replica partition function, Eq.(15), we obtain the standard expression

$$Z(N, L) = \prod_{a=1}^N \int_{\phi_a(0)=0}^{\phi_a(L)=0} \mathcal{D}\phi_a(\tau) e^{-\beta H_N[\phi]} \quad (17)$$

where $\phi \equiv \{\phi_1, \dots, \phi_N\}$. According to the above definition this partition function describe the statistics of N δ -interacting (attracting) trajectories $\phi_a(\tau)$ all starting (at $\tau = 0$) and ending (at $\tau = L$) at zero: $\phi_a(0) = \phi_a(L) = 0$

In order to map the problem to one-dimensional quantum bosons, let us introduce more general object

$$\Psi(\mathbf{x}; t) = \prod_{a=1}^N \int_{\phi_a(0)=0}^{\phi_a(t)=x_a} \mathcal{D}\phi_a(\tau) e^{-\beta H_N[\phi]} \quad (18)$$

which describes N trajectories $\phi_a(\tau)$ all starting at zero ($\phi_a(0) = 0$), but ending at $\tau = t$ in arbitrary given points $\{x_1, \dots, x_N\}$. One can easily show that instead of using the path integral, $\Psi(\mathbf{x}; t)$ may be obtained as the solution of the linear differential equation

$$\partial_t \Psi(\mathbf{x}; t) = \frac{1}{2\beta} \sum_{a=1}^N \partial_{x_a}^2 \Psi(\mathbf{x}; t) + \frac{1}{2} \beta^2 u \sum_{a \neq b}^N \delta(x_a - x_b) \Psi(\mathbf{x}; t) \quad (19)$$

with the initial condition

$$\Psi(\mathbf{x}; 0) = \prod_{a=1}^N \delta(x_a) \quad (20)$$

One can easily see that Eq.(19) is the imaginary-time Schrödinger equation

$$-\partial_t \Psi(\mathbf{x}; t) = \hat{H} \Psi(\mathbf{x}; t) \quad (21)$$

with the Hamiltonian

$$\hat{H} = -\frac{1}{2\beta} \sum_{a=1}^N \partial_{x_a}^2 - \frac{1}{2} \beta^2 u \sum_{a \neq b}^N \delta(x_a - x_b) \quad (22)$$

which describes N bose-particles of mass β interacting via the *attractive* two-body potential $-\beta^2 u \delta(x)$. The original replica partition function, Eq.(17), then is obtained via a particular choice of the final-point coordinates,

$$Z(N, L) = \Psi(\mathbf{0}; L). \quad (23)$$

The eigenfunction of N -particle system of *repulsive* bosons ($u < 0$), Eq.(22), have been derived by Lieb and Liniger in 1963 [28] (for details see Appendix A, as well as Refs. [29, 30]). The spectrum and some properties of the eigenfunctions for attractive ($u > 0$) one-dimensional quantum boson system have been derived by McGuire [31] and

by Yang [32] (see also Ref. [33, 34]). Detailed structure and the properties of these wave functions are described in Appendix B. A generic eigenstate of such system consists of M ($1 \leq M \leq N$) "clusters" $\{\Omega_\alpha\}$ ($\alpha = 1, \dots, M$) of bound particles. Each cluster is characterized by the momentum q_α of its center of mass motion, and by the number n_α of particles contained in it (such that $\sum_{\alpha=1}^M n_\alpha = N$). Correspondingly, the eigenfunction $\Psi_{\mathbf{q}, \mathbf{n}}^{(M)}(x_1, \dots, x_N)$ of such state is characterized by M continuous parameters $\mathbf{q} = (q_1, \dots, q_M)$ and M integer parameters $\mathbf{n} = (n_1, \dots, n_M)$ (see Appendix B2, Eq.(B.28)). The energy spectrum of this state is

$$E_M(\mathbf{q}, \mathbf{n}) = \frac{1}{2\beta} \sum_{\alpha=1}^M n_\alpha q_\alpha^2 - \frac{\kappa^2}{24\beta} \sum_{\alpha=1}^M (n_\alpha^3 - n_\alpha) \quad (24)$$

where

$$\kappa = \beta^3 u \quad (25)$$

A general time dependent solution $\Psi(\mathbf{x}, t)$ of the Schrödinger equation (19) with the initial conditions, Eq.(20), can be represented in the form of the linear combination of the eigenfunctions $\Psi_{\mathbf{q}, \mathbf{n}}^{(M)}(\mathbf{x})$ (Appendix B4, Eq.(B.51)):

$$\Psi(\mathbf{x}, t) = \sum_{M=1}^N \sum_{\mathbf{n}}' \int' \mathcal{D}\mathbf{q} \Psi_{\mathbf{q}, \mathbf{n}}^{(M)}(\mathbf{x}) \Psi_{\mathbf{q}, \mathbf{n}}^{(M)*}(\mathbf{0}) \exp[-E_M(\mathbf{q}, \mathbf{n}) t] \quad (26)$$

where the summations over the integer parameters n_α and the integrations over the momenta q_α are performed in a restricted subspace, Eqs.(B.43)-(B.46) and (B.52), which reflects the specific symmetry properties of the eigenfunctions $\Psi_{\mathbf{q}, \mathbf{n}}^{(M)}(\mathbf{x})$. Correspondingly, according to Eq.(23), for the replica partition function of the original directed polymer problem one gets (Appendix B4, Eq.(B.60))

$$Z(N, L) = \sum_{M=1}^N \frac{1}{M!} \left[\prod_{\alpha=1}^M \int_{-\infty}^{+\infty} \frac{dq_\alpha}{2\pi} \sum_{n_\alpha=1}^{\infty} \right] \delta\left(\sum_{\alpha=1}^M n_\alpha, N\right) |\Psi_{\mathbf{q}, \mathbf{n}}^{(M)}(\mathbf{0})|^2 e^{-E_M(\mathbf{q}, \mathbf{n})L} \quad (27)$$

where due to the symmetry of the function $f(\mathbf{q}, \mathbf{n}) = |\Psi_{\mathbf{q}, \mathbf{n}}^{(M)}(\mathbf{0})|^2 \exp[-E_M(\mathbf{q}, \mathbf{n})L]$ with respect to permutations of all its M pairs of arguments (q_α, n_α) the integrations over M momenta q_α can be extended to the whole space R_M while the summations over n_α 's are bounded by the only constrain $\sum_{\alpha=1}^M n_\alpha = N$ (for simplicity, due to the presence of the Kronecker symbol $\delta(\sum_{\alpha=1}^M n_\alpha, N)$, the summations over n_α 's are extended to infinity).

III. REPLICAS PARTITION FUNCTION AND THE FREE ENERGY DISTRIBUTION FUNCTION

Using the explicit form of the wave functions $\Psi_{\mathbf{q}, \mathbf{n}}^{(M)}(\mathbf{x})$, Eq.(B.28), the expression in Eq.(27) for the replica partition function can be reduced to (Appendix B4, Eq.(B.62)-(B.63))

$$Z(N, L) = e^{-\beta N L f_0} \tilde{Z}(N, \lambda) \quad (28)$$

where $f_0 = \frac{1}{24}\beta^4 u^2 - \frac{1}{\beta L} \ln(\beta^3 u)$ is the linear (selfaveraging) free energy density (cf. Eq.(7), and

$$\begin{aligned} \tilde{Z}(N, L) = & N! \left\{ \int_{-\infty}^{+\infty} \frac{dq}{2\pi\kappa N} \exp\left[-\frac{L}{2\beta} N q^2 + \frac{\kappa^2 L}{24\beta} N^3\right] + \right. \\ & + \sum_{M=2}^N \frac{1}{M!} \left[\prod_{\alpha=1}^M \sum_{n_\alpha=1}^{\infty} \int_{-\infty}^{+\infty} \frac{dq_\alpha}{2\pi\kappa n_\alpha} \right] \delta\left(\sum_{\alpha=1}^M n_\alpha, N\right) \prod_{\alpha < \beta}^M \frac{|q_\alpha - q_\beta - \frac{i\kappa}{2}(n_\alpha - n_\beta)|^2}{|q_\alpha - q_\beta - \frac{i\kappa}{2}(n_\alpha + n_\beta)|^2} \times \\ & \times \exp\left[-\frac{L}{2\beta} \sum_{\alpha=1}^M n_\alpha q_\alpha^2 + \frac{\kappa^2 L}{24\beta} \sum_{\alpha=1}^M n_\alpha^3\right] \Big\} \quad (29) \end{aligned}$$

The first term in the above expression is the contribution of the ground state ($M = 1$), while the next terms ($M \geq 2$) are the contributions of the rest of the energy spectrum.

The terms cubic in n_α in the exponential of Eq. (29) can be linearised with the help of Airy function, using the standard relation

$$\exp\left(\frac{1}{3}\lambda^3 n^3\right) = \int_{-\infty}^{+\infty} dy \text{Ai}(y) \exp(\lambda n) \quad (30)$$

Redefining the momenta, $q_\alpha = (\beta\kappa/L)^{1/3} p_\alpha$ and introducing a new parameter

$$\lambda(L) = \frac{1}{2} \left(\frac{L}{\beta} \kappa^2 \right)^{1/3} = \frac{1}{2} (\beta^5 u^2 L)^{1/3} \quad (31)$$

after shifting the Airy function parameters of integration $y_\alpha \rightarrow y_\alpha + p_\alpha^2$ the expression for $\tilde{Z}(N, \lambda)$ becomes sufficiently compact:

$$\begin{aligned} \tilde{Z}(N, \lambda) = & N! \int \int_{-\infty}^{+\infty} \frac{dy dp}{4\pi\lambda N} \text{Ai}(y + p^2) e^{\lambda N y} + \\ & + N! \sum_{M=2}^N \frac{1}{M!} \left[\prod_{\alpha=1}^M \sum_{n_\alpha=1}^{\infty} \int \int_{-\infty}^{+\infty} \frac{dy_\alpha dp_\alpha}{4\pi\lambda n_\alpha} \text{Ai}(y_\alpha + p_\alpha^2) e^{\lambda n_\alpha y_\alpha} \right] \prod_{\alpha < \beta}^M \frac{|\lambda(n_\alpha - n_\beta) - i(p_\alpha - p_\beta)|^2}{|\lambda(n_\alpha + n_\beta) - i(p_\alpha - p_\beta)|^2} \delta\left(\sum_{\alpha=1}^M n_\alpha, N\right) \end{aligned} \quad (32)$$

Now, using the Cauchy double alternant identity

$$\frac{\prod_{\alpha < \beta}^M (a_\alpha - a_\beta)(b_\alpha - b_\beta)}{\prod_{\alpha, \beta=1}^M (a_\alpha - b_\beta)} = (-1)^{M(M-1)/2} \det \left[\frac{1}{a_\alpha - b_\beta} \right]_{\alpha, \beta=1, \dots, M} \quad (33)$$

the product term in eq.(32) can be represented in the determinant form:

$$\prod_{\alpha < \beta}^M \frac{|\lambda(n_\alpha - n_\beta) - i(p_\alpha - p_\beta)|^2}{|\lambda(n_\alpha + n_\beta) - i(p_\alpha - p_\beta)|^2} = \left[\prod_{\alpha=1}^M (2\lambda n_\alpha) \right] \det \left[\frac{1}{\lambda n_\alpha - ip_\alpha + \lambda n_\beta + ip_\beta} \right]_{\alpha, \beta=1, \dots, M} \quad (34)$$

Substituting now the expression for the replica partition function $\tilde{Z}(N, \lambda)$ into the definition of the probability function, eq.(12), we can perform summation over N (which would lift the constraint $\sum_{\alpha=1}^M n_\alpha = N$) and obtain:

$$W(x) = \lim_{\lambda \rightarrow \infty} \left\{ 1 + \sum_{M=1}^{\infty} \frac{(-1)^M}{M!} \left[\prod_{\alpha=1}^M \int \int_{-\infty}^{+\infty} \frac{dy_\alpha dp_\alpha}{2\pi} \text{Ai}(y_\alpha + p_\alpha^2) \sum_{n_\alpha=1}^{\infty} (-1)^{n_\alpha-1} e^{\lambda n_\alpha (y_\alpha + x)} \right] \det \left[\frac{1}{\lambda n_\alpha - ip_\alpha + \lambda n_\beta + ip_\beta} \right] \right\} \quad (35)$$

The above expression is nothing else but the expansion of the Fredholm determinant $\det(1 - \hat{K})$ (see e.g. [35]) with the kernel

$$\hat{K} \equiv K[(n, p); (n', p')] = \left[\int_{-\infty}^{+\infty} dy \text{Ai}(y + p^2) (-1)^{n-1} e^{\lambda n (y+x)} \right] \frac{1}{\lambda n - ip + \lambda n' + ip'} \quad (36)$$

Using the exponential representation of this determinant we get

$$W(x) = \lim_{\lambda \rightarrow \infty} \exp \left[- \sum_{M=1}^{\infty} \frac{1}{M} \text{Tr} \hat{K}^M \right] \quad (37)$$

where

$$\begin{aligned} \text{Tr} \hat{K}^M = & \left[\prod_{\alpha=1}^M \int \int_{-\infty}^{+\infty} \frac{dy_\alpha dp_\alpha}{2\pi} \text{Ai}(y_\alpha + p_\alpha^2) \sum_{n_\alpha=1}^{\infty} (-1)^{n_\alpha-1} e^{\lambda n_\alpha (y_\alpha + x)} \right] \times \\ & \times \frac{1}{(\lambda n_1 - ip_1 + \lambda n_2 + ip_2)(\lambda n_2 - ip_2 + \lambda n_3 + ip_3) \dots (\lambda n_M - ip_M + \lambda n_1 + ip_1)} \end{aligned} \quad (38)$$

Substituting here

$$\frac{1}{\lambda n_\alpha - ip_\alpha + \lambda n_{\alpha+1} + ip_{\alpha+1}} = \int_0^\infty d\omega_\alpha \exp[-(\lambda n_\alpha - ip_\alpha + \lambda n_{\alpha+1} + ip_{\alpha+1})\omega_\alpha] \quad (39)$$

one can easily perform the summation over n_α 's. Taking into account that

$$\lim_{\lambda \rightarrow \infty} \sum_{n=1}^{\infty} (-1)^{n-1} e^{\lambda n z} = \lim_{\lambda \rightarrow \infty} \frac{e^{\lambda z}}{1 + e^{\lambda z}} = \theta(z) \quad (40)$$

and shifting the integration parameters, $y_\alpha \rightarrow y_\alpha - x + \omega_\alpha + \omega_{\alpha-1}$ and $\omega_\alpha \rightarrow \omega_\alpha + x/2$, after taking the thermodynamic limit, $L \rightarrow \infty$ (which according to Eq.(31) is equivalent to $\lambda \rightarrow \infty$) we obtain

$$\lim_{\lambda \rightarrow \infty} Tr \hat{K}^M = \prod_{\alpha=1}^M \int_0^\infty dy_\alpha \int_{-\infty}^{+\infty} \frac{dp_\alpha}{2\pi} \int_{-x/2}^\infty d\omega_\alpha Ai(y_\alpha + p_\alpha^2 + \omega_\alpha + \omega_{\alpha-1}) e^{ip_\alpha(\omega_\alpha - \omega_{\alpha-1})} \quad (41)$$

where by definition it is assumed that $\omega_0 \equiv \omega_M$. Using the Airy function integral representation, and taking into account that it satisfies the differential equation, $Ai''(t) = t Ai(t)$, one can easily perform the following integrations:

$$\begin{aligned} \int_0^\infty dy \int_{-\infty}^{+\infty} \frac{dp}{2\pi} Ai(y + p^2 + \omega + \omega') e^{ip(\omega - \omega')} &= 2^{-1/3} \int_0^\infty dy Ai(2^{1/3}\omega + y) Ai(2^{1/3}\omega' + y) \\ &= \frac{Ai(2^{1/3}\omega) Ai'(2^{1/3}\omega') - Ai'(2^{1/3}\omega) Ai(2^{1/3}\omega')}{\omega - \omega'} \end{aligned} \quad (42)$$

Redefining $\omega_\alpha \rightarrow \omega_\alpha 2^{-1/3}$ we find

$$\lim_{\lambda \rightarrow \infty} Tr \hat{K}^M = \int \int \dots \int_{-x/2^{2/3}}^\infty d\omega_1 d\omega_2 \dots d\omega_M K_A(\omega_1, \omega_2) K_A(\omega_2, \omega_3) \dots K_A(\omega_M, \omega_1) \quad (43)$$

where

$$K_A(\omega, \omega') = \frac{Ai(\omega) Ai'(\omega') - Ai'(\omega) Ai(\omega')}{\omega - \omega'} \quad (44)$$

is the so called Airy kernel. This proves that in the thermodynamic limit, $L \rightarrow \infty$, the probability function $W(x)$, eq.(10), is defined by the Fredholm determinant,

$$W(x) = \det[1 - \hat{K}_A] \equiv F_2(-x/2^{2/3}) \quad (45)$$

where $\hat{K}_A$ is the integral operator on $[-x/2^{2/3}, \infty)$ with the Airy kernel, eq.(44). The function $F_2(s)$ is the Tracy-Widom distribution [14]

$$F_2(s) = \exp\left(-\int_s^\infty dt (s-t) q^2(t)\right) \quad (46)$$

where the function $q(t)$ is the solution of the Painlevé II equation, $q'' = tq + 2q^3$ with the boundary condition, $q(t \rightarrow +\infty) \sim Ai(t)$ (see Appendix C).

IV. CONCLUSIONS

The Tracy-Widom distribution, Eq.(46), was originally derived as the probability distribution of the largest eigenvalue of a $n \times n$ random hermitian matrix in the limit $n \rightarrow \infty$. At present there exists an appreciable list of statistical systems (which are not always look similar) in which the fluctuations of the quantities which play the role of "energy" are described by *the same* distribution function $F_2(s)$. These systems are: the polynuclear growth (PNG) model [15], the longest increasing subsequences (LIS) model [16], the longest common subsequences (LCS) [17], the oriented digital boiling model [18], the ballistic decomposition model [19], the zero-temperature lattice version of the directed polymers with an exponential and geometric site-disorder distribution [20]. Now we can add to this list the continuous one-dimensional directed polymers with Gaussian δ -correlated random potential.

Appendix A

Quantum bosons with repulsive interactions

1. Eigenfunctions

The eigenstates equation for N -particle system of one-dimensional quantum bosons with δ -interactions is

$$\frac{1}{2} \sum_{a=1}^N \partial_{x_a}^2 \Psi(\mathbf{x}) + \frac{1}{2} \kappa \sum_{a \neq b}^N \delta(x_a - x_b) \Psi(\mathbf{x}) = -\beta E \Psi(\mathbf{x}) \quad (\text{A.1})$$

(where $\kappa = \beta^3 u$). Due to the symmetry of the wave function with respect to permutations of its arguments it is sufficient to consider it in the sector

$$x_1 < x_2 < \dots < x_N \quad (\text{A.2})$$

as well as at its boundary. Inside this sector the wave function $\Psi(\mathbf{x})$ satisfy the equation

$$\frac{1}{2} \sum_{a=1}^N \partial_{x_a}^2 \Psi(\mathbf{x}) = -\beta E \Psi(\mathbf{x}) \quad (\text{A.3})$$

which describes N free particles, and its generic solution is the linear combination of N plane waves characterized by N momenta $\{q_1, q_2, \dots, q_N\} \equiv \mathbf{q}$. Integrating Eq.(A.1) over the variable $(x_{i+1} - x_i)$ in a small interval around zero, $|x_{i+1} - x_i| < \epsilon \rightarrow 0$, and assuming that the other variables $\{x_j\}$ (with $j \neq i, i+1$) belong to the sector, Eq.(A.2), one easily finds that the wave function $\Psi(\mathbf{x})$ must satisfy the following boundary conditions:

$$(\partial_{x_{i+1}} - \partial_{x_i} + \kappa) \Psi(\mathbf{x}) \Big|_{x_{i+1}=x_i+0} = 0 \quad (\text{A.4})$$

Functions satisfying both Eq. (A.3) and the boundary conditions Eq. (A.4) can be written in the form

$$\Psi_{q_1 \dots q_N}(x_1, \dots, x_N) \equiv \Psi_{\mathbf{q}}^{(N)}(\mathbf{x}) = C \left(\prod_{a < b}^N [\partial_{x_a} - \partial_{x_b} + \kappa] \right) \det \left[\exp(iq_c x_d) \right]_{(c,d)=1, \dots, N} \quad (\text{A.5})$$

where C is the normalization constant to be defined later. First of all, it is evident that being the linear combination of the plane waves, the above wave function satisfy Eq.(A.3). To demonstrate which way this function satisfy the boundary conditions, Eq.(A.4), let us check it, as an example, for the case $i = 1$. According to Eq.(A.5), the wave function $\Psi_{\mathbf{q}}^{(N)}(\mathbf{x})$ can be represented in the form

$$\Psi_{\mathbf{q}}^{(N)}(\mathbf{x}) = -(\partial_{x_2} - \partial_{x_1} - \kappa) \tilde{\Psi}_{\mathbf{q}}^{(N)}(\mathbf{x}) \quad (\text{A.6})$$

where

$$\tilde{\Psi}_{\mathbf{q}}^{(N)}(\mathbf{x}) = C \left(\prod_{a=3}^N [\partial_{x_1} - \partial_{x_a} + \kappa] [\partial_{x_2} - \partial_{x_a} + \kappa] \right) \left(\prod_{3 \leq a < b}^N [\partial_{x_a} - \partial_{x_b} + \kappa] \right) \det \left[\exp(iq_c x_d) \right]_{(c,d)=1, \dots, N} \quad (\text{A.7})$$

One can easily see that this function is *antisymmetric* with respect to the permutation of x_1 and x_2 . Substituting Eq.(A.6) into Eq.(A.4) (with $i = 1$) we get

$$-\left[(\partial_{x_2} - \partial_{x_1})^2 - \kappa^2 \right] \tilde{\Psi}_{\mathbf{q}}^{(N)}(\mathbf{x}) \Big|_{x_2=x_1} = 0 \quad (\text{A.8})$$

Given the antisymmetry of the l.h.s expression with respect to the permutation of x_1 and x_2 the above condition is indeed satisfied at boundary $x_1 = x_2$.

Since the eigenfunction $\Psi_{\mathbf{q}}^{(N)}(\mathbf{x})$ satisfying Eq.(A.1) must be *symmetric* with respect to permutations of its arguments, the function, Eq.(A.5), can be easily continued beyond the sector, Eq.(A.2), to the entire space of variables $\{x_1, x_2, \dots, x_N\} \in R_N$,

$$\Psi_{\mathbf{q}}^{(N)}(\mathbf{x}) = C \left(\prod_{a < b}^N \left[-i(\partial_{x_a} - \partial_{x_b}) + i\kappa \operatorname{sgn}(x_a - x_b) \right] \right) \det \left[\exp(iq_c x_d) \right]_{(c,d)=1, \dots, N} \quad (\text{A.9})$$

where, by definition, the differential operators ∂_{x_a} act only on the exponential terms and not on the $\operatorname{sgn}(x)$ functions, and for further convenience we have redefined $i^{N(N-1)/2} C \rightarrow C$. Explicitly the determinant in the above equation is

$$\det \left[\exp(iq_c x_d) \right]_{(c,d)=1, \dots, N} = \sum_P (-1)^{[P]} \exp \left[i \sum_{a=1}^N q_{p_a} x_a \right] \quad (\text{A.10})$$

where the summation goes over the permutations P of N momenta $\{q_1, q_2, \dots, q_N\}$ over N particles $\{x_1, x_2, \dots, x_N\}$, and $[P]$ denotes the parity of the permutation. In this way the eigenfunction, Eq.(A.9), can be represented as follows

$$\Psi_{\mathbf{q}}^{(N)}(\mathbf{x}) = C \sum_P (-1)^{[P]} \left(\prod_{a < b}^N \left[-i(\partial_{x_a} - \partial_{x_b}) + i\kappa \operatorname{sgn}(x_a - x_b) \right] \right) \exp \left[i \sum_{a=1}^N q_{p_a} x_a \right] \quad (\text{A.11})$$

Taking the derivatives, we obtain

$$\Psi_{\mathbf{q}}^{(N)}(\mathbf{x}) = C \sum_P (-1)^{[P]} \left(\prod_{a < b}^N \left[q_{p_a} - q_{p_b} + i\kappa \operatorname{sgn}(x_a - x_b) \right] \right) \exp \left[i \sum_{a=1}^N q_{p_a} x_a \right] \quad (\text{A.12})$$

It is evident from these representations that the eigenfunctions $\Psi_{\mathbf{q}}^{(N)}(\mathbf{x})$ are *antisymmetric* with respect to permutations of the momenta $q_1, \dots, q_N$.

Finally, substituting the expression for the eigenfunctions, Eq.(A.5) (which is valid in the sector, Eq.(A.2)), into Eq.(A.3) for the energy spectrum we find

$$E = \frac{1}{2\beta} \sum_{a=1}^N q_a^2 \quad (\text{A.13})$$

2. Orthonormality

Now one can easily prove that the above eigenfunctions with different momenta are orthogonal to each other. Let us consider two wave functions $\Psi_{\mathbf{q}}^{(N)}(\mathbf{x})$ and $\Psi_{\mathbf{q}'}^{(N)}(\mathbf{x})$ where it is assumed that

$$\begin{aligned} q_1 &< q_2 < \dots < q_N \\ q'_1 &< q'_2 < \dots < q'_N \end{aligned} \quad (\text{A.14})$$

Using the representation, Eq.(A.11), for the overlap of these two function we get

$$\begin{aligned} \overline{\Psi_{\mathbf{q}'}^{(N)*}(\mathbf{x}) \Psi_{\mathbf{q}}^{(N)}(\mathbf{x})} &\equiv \int_{-\infty}^{+\infty} d^N \mathbf{x} \Psi_{\mathbf{q}'}^{(N)*}(\mathbf{x}) \Psi_{\mathbf{q}}^{(N)}(\mathbf{x}) \\ &= |C|^2 \sum_{P, P'} (-1)^{[P]+[P']} \int_{-\infty}^{+\infty} d^N \mathbf{x} \left\{ \left(\prod_{a < b}^N \left[i(\partial_{x_a} - \partial_{x_b}) - i\kappa \operatorname{sgn}(x_a - x_b) \right] \right) \exp \left[-i \sum_{a=1}^N q'_{p'_a} x_a \right] \right\} \times \\ &\times \left\{ \left(\prod_{a < b}^N \left[-i(\partial_{x_a} - \partial_{x_b}) + i\kappa \operatorname{sgn}(x_a - x_b) \right] \right) \exp \left[i \sum_{a=1}^N q_{p_a} x_a \right] \right\} \end{aligned} \quad (\text{A.15})$$

Integrating by parts we obtain

$$\begin{aligned} \overline{\Psi_{\mathbf{q}'}^{(N)*}(\mathbf{x})\Psi_{\mathbf{q}}^{(N)}(\mathbf{x})} &= |C|^2 \sum_{P,P'} (-1)^{[P]+[P']} \int_{-\infty}^{+\infty} d^N \mathbf{x} \exp\left[-i \sum_{a=1}^N q'_{p'_a} x_a\right] \times \\ &\times \left(\prod_{a<b}^N [-i(\partial_{x_a} - \partial_{x_b}) - i\kappa \operatorname{sgn}(x_a - x_b)] [-i(\partial_{x_a} - \partial_{x_b}) + i\kappa \operatorname{sgn}(x_a - x_b)] \right) \exp\left[i \sum_{a=1}^N q_{p_a} x_a\right] \end{aligned} \quad (\text{A.16})$$

or

$$\overline{\Psi_{\mathbf{q}'}^{(N)*}(\mathbf{x})\Psi_{\mathbf{q}}^{(N)}(\mathbf{x})} = |C|^2 \sum_{P,P'} (-1)^{[P]+[P']} \int_{-\infty}^{+\infty} d^N \mathbf{x} \exp\left[-i \sum_{a=1}^N q'_{p'_a} x_a\right] \left(\prod_{a<b}^N [-(\partial_{x_a} - \partial_{x_b})^2 + \kappa^2] \right) \exp\left[i \sum_{a=1}^N q_{p_a} x_a\right] \quad (\text{A.17})$$

Taking the derivatives and performing the integrations we find

$$\begin{aligned} \overline{\Psi_{\mathbf{q}'}^{(N)*}(\mathbf{x})\Psi_{\mathbf{q}}^{(N)}(\mathbf{x})} &= |C|^2 \sum_{P,P'} (-1)^{[P]+[P']} \left(\prod_{a<b}^N [(q_{p_a} - q_{p_b})^2 + \kappa^2] \right) \int_{-\infty}^{+\infty} d^N \mathbf{x} \exp\left[i \sum_{a=1}^N (q_{p_a} - q'_{p'_a}) x_a\right] \\ &= |C|^2 \sum_{P,P'} (-1)^{[P]+[P']} \left(\prod_{a<b}^N [(q_{p_a} - q_{p_b})^2 + \kappa^2] \right) \left[\prod_{a=1}^N (2\pi) \delta(q_{p_a} - q'_{p'_a}) \right] \end{aligned} \quad (\text{A.18})$$

Taking into account the constraint, Eq.(A.14), one can easily note that the only the terms which survive in the above summation over the permutations are $P = P'$, all contributing equal value. Thus, we finally get

$$\overline{\Psi_{\mathbf{q}'}^{(N)*}(\mathbf{x})\Psi_{\mathbf{q}}^{(N)}(\mathbf{x})} = |C|^2 N! \left(\prod_{a<b}^N [(q_a - q_b)^2 + \kappa^2] \right) \left[\prod_{a=1}^N (2\pi) \delta(q_a - q'_a) \right] \quad (\text{A.19})$$

With the normalization constant

$$|C(\mathbf{q})|^2 = \frac{1}{N! \prod_{a<b}^N [(q_a - q_b)^2 + \kappa^2]} \quad (\text{A.20})$$

we conclude that the set of the eigenfunctions, Eq.(A.11) or (A.12), are orthonormal. The proof of completeness of this set is given in Ref. [30]. It should be noted that the above wave functions present the orthonormal set of eigenfunctions of the problem, Eq.(A.1), for any sign of the interactions κ , e.i. both for the repulsive, $\kappa < 0$, and for the attractive, $\kappa > 0$, cases. However, only in the case of repulsion this set is complete, while in the case of attractive interactions, $\kappa > 0$, in addition to the solutions, Eq.(A.11), which describe the continuous free particles spectrum, one finds the whole family of discrete bound eigenstates (which do not exist in the case of repulsion).

Appendix B

Quantum bosons with attractive interactions

1. Ground state

The simplest example of the bound eigenstate defined by eq.(A.1) (with $\kappa > 0$) is the one in which all N particles are bound into a single "cluster":

$$\Psi_q^{(1)}(\mathbf{x}) = C \exp\left[iq \sum_{a=1}^N x_a - \frac{1}{4}\kappa \sum_{a,b=1}^N |x_a - x_b|\right] \quad (\text{B.1})$$

where C is the normalization constant (to be defined below) and q is the continuous momentum of free center of mass motion. Substituting this function in Eq.(A.1), one can easily check that this is indeed the eigenfunction with the energy spectrum given by the relation

$$E = -\frac{1}{2\beta} \sum_{a=1}^N \left[iq - \frac{1}{2} \kappa \sum_{b=1}^N \text{sgn}(x_a - x_b) \right]^2 \quad (\text{B.2})$$

where it is assumed (by definition) that $\text{sgn}(0) = 0$. Since the result of the above summations does not depend on the mutual particles positions, for simplicity we can order them according to Eq.(A.2). Then, using well known relations

$$\sum_{b=1}^N \text{sgn}(x_a - x_b) = -(N + 1 - 2a) \quad (\text{B.3})$$

$$\sum_{a=1}^N a = \frac{1}{2} N(N + 1) \quad (\text{B.4})$$

$$\sum_{a=1}^N a^2 = \frac{1}{6} (N + 1)(2N + 1) \quad (\text{B.5})$$

for the energy spectrum, Eq.(B.2), we get

$$E = \frac{N}{2\beta} q^2 - \frac{\kappa^2}{24\beta} (N^3 - N) \equiv E_1(q, N) \quad (\text{B.6})$$

The normalization constant C is defined by the orthonormality condition

$$\overline{\Psi_{q'}^{(1)*}(\mathbf{x}) \Psi_q^{(1)}(\mathbf{x})} \equiv \int_{-\infty}^{+\infty} dx_1 \dots dx_N \Psi_{q'}^{(1)*}(\mathbf{x}) \Psi_q^{(1)}(\mathbf{x}) = (2\pi) \delta(q - q') \quad (\text{B.7})$$

Substituting here Eq.(B.1) we get

$$\begin{aligned} \overline{\Psi_{q'}^{(1)*}(\mathbf{x}) \Psi_q^{(1)}(\mathbf{x})} &= |C|^2 \int_{-\infty}^{+\infty} dx_1 \dots dx_N \exp \left[i(q - q') \sum_{a=1}^N x_a - \frac{1}{2} \kappa \sum_{a,b=1}^N |x_a - x_b| \right] \\ &= |C|^2 N! \int_{-\infty}^{+\infty} dx_1 \int_{x_1}^{+\infty} dx_2 \dots \int_{x_{N-1}}^{+\infty} dx_N \exp \left[i(q - q') \sum_{a=1}^N x_a + \kappa \sum_{a=1}^N (N + 1 - 2a) x_a \right] \end{aligned} \quad (\text{B.8})$$

where for the ordering, Eq.(A.2), we have used the relation

$$\frac{1}{2} \sum_{a,b=1}^N |x_a - x_b| = - \sum_{a=1}^N (N + 1 - 2a) x_a \quad (\text{B.9})$$

Integrating first over x_N , then over x_{N-1} , and proceeding until x_1 , we find

$$\begin{aligned} \overline{\Psi_{q'}^{(1)*}(\mathbf{x}) \Psi_q^{(1)}(\mathbf{x})} &= |C|^2 N! \left(\prod_{r=1}^{N-1} \frac{1}{r[(N-r)\kappa - i(q - q')]} \right) \int_{-\infty}^{+\infty} dx_1 \exp [iN(q - q')x_1] \\ &= |C|^2 N! \left(\prod_{r=1}^{N-1} \frac{1}{r(N-r)\kappa} \right) (2\pi) \delta(N(q - q')) \\ &= |C|^2 \frac{N\kappa}{N! \kappa^N} (2\pi) \delta(q - q') \end{aligned} \quad (\text{B.10})$$

According to Eq.(B.7) this defines the normalization constant

$$C = \sqrt{\frac{\kappa^N N!}{\kappa N}} \equiv C^{(1)}(q) \quad (\text{B.11})$$

Note that the eigenstate described by the considered wave function, Eq.(B.1), exists only in the case of attraction, $\kappa > 0$, otherwise this function is divergent at infinity and consequently it is not normalizable.

It should be noted that the wave function, Eq.(B.1), can also be derived from the general eigenfunctions structure, Eq.(A.12), by introducing (discrete) imaginary parts for the momenta q_a . We assume again that the position of particles are ordered according to Eq.(A.2), and define the particles' momenta according to the rule

$$q_a = q - \frac{i}{2}\kappa(N+1-2a) \quad (\text{B.12})$$

Substituting this into Eq.(A.12) we get

$$\begin{aligned} \Psi_q^{(1)}(x_1 < x_2 < \dots < x_N) &\propto \sum_P (-1)^{[P]} \left(\prod_{a < b}^N \left[\left(q - \frac{i}{2}\kappa(N+1-2P_a) \right) - \left(q - \frac{i}{2}\kappa(N+1-2P_b) \right) - i\kappa \right] \right) \times \\ &\times \exp \left[iq \sum_{a=1}^N x_a + \frac{\kappa}{2} \sum_{a=1}^N (N+1-2P_a)x_a \right] \end{aligned} \quad (\text{B.13})$$

$$\propto \sum_P (-1)^{[P]} \left(\prod_{a < b}^N [P_b - P_a + 1] \right) \exp \left[iq \sum_{a=1}^N x_a + \frac{\kappa}{2} \sum_{a=1}^N (N+1-2P_a)x_a \right] \quad (\text{B.14})$$

Here one can easily note that due to the presence of the product $\prod_{a < b}^N [P_b - P_a + 1]$ in the summation over permutations only the trivial one, $P_a = a$, gives non-zero contribution (if we permute any two numbers in the sequence $1, 2, \dots, N$ then we can always find two numbers $a < b$, such that $P_b = P_a - 1$). Thus

$$\Psi_q^{(1)}(x_1 < x_2 < \dots < x_N) \propto \exp \left[iq \sum_{a=1}^N x_a + \frac{\kappa}{2} \sum_{a=1}^N (N+1-2a)x_a \right] \quad (\text{B.15})$$

Taking into account the relation, Eq.(B.9), we recover the function, Eq.(B.1), which is symmetric with respect to its N arguments and therefore can be extended beyond the sector, Eq.(A.2), for arbitrary particles positions. Finally, substituting the momenta, Eq.(B.12), into the general expression for the energy spectrum, Eq.(A.13), we get

$$E = \frac{1}{2\beta} \sum_{a=1}^N \left[q - \frac{i}{2}\kappa(N+1-2a) \right]^2 \quad (\text{B.16})$$

Performing here simple summations (using Eqs.(B.4), (B.5)) one recovers Eq.(B.6).

2. Eigenfunctions

A generic eigenfunction of attractive bosons is characterized by N momenta parameters $\{q_a\}$ ($a = 1, 2, \dots, N$) which may have imaginary parts. It is convenient to group these parameters into M ($1 \leq M \leq N$) "vector" momenta,

$$q_r^\alpha = q_\alpha - \frac{i}{2}\kappa(n_\alpha + 1 - 2r) \quad (\text{B.17})$$

where q_α ($\alpha = 1, 2, \dots, M$) are the continuous (real) parameters, and the discrete imaginary components of each "vector" are labeled by an index $r = 1, 2, \dots, n_\alpha$. With the given total number of particles equal to N , the integers n_α have to satisfy the constraint

$$\sum_{\alpha=1}^M n_\alpha = N \quad (\text{B.18})$$

In other words, a generic eigenstate is characterized by the discrete number M of complex "vector" momenta, by the set of M integer parameters $\{n_1, n_2, \dots, n_M\} \equiv \mathbf{n}$ (which are the numbers of imaginary components of each "vector") and by the set of M real continuous momenta $\{q_1, q_2, \dots, q_M\} \equiv \mathbf{q}$.

The general expression for the eigenfunctions is given in Eqs.(A.9)-(A.12). To understand the structure of the determinant of the $N \times N$ matrix $\exp(iq_a x_b)$, which defines these wave functions, the N momenta q_a , eq.(B.17), can be ordered as follows:

$$\{q_a\} \equiv \{q_r^\alpha\} = \{q_1^1, q_2^1, \dots, q_{n_1}^1; q_1^2, q_2^2, \dots, q_{n_2}^2; \dots; q_1^M, q_2^M, \dots, q_{n_M}^M\} \quad (\text{B.19})$$

By definition,

$$\det \left[\exp(iq_a x_c) \right]_{(c,d)=1,\dots,N} = \sum_P (-1)^{[P]} \exp \left[i \sum_{a=1}^N q_{p_a} x_a \right] \quad (\text{B.20})$$

where the summation goes over the permutations of N momenta $\{q_a\}$, Eq.(B.19), over N particles $\{x_1, x_2, \dots, x_N\}$, and $[P]$ denotes the parity of the permutation. For a given permutation P a particle number a is attributed a momentum component $q_{r(a)}^{\alpha(a)}$. The particles getting the momenta with the same α (having the same real part q_α) will be called belonging to a cluster Ω_α . For a given permutation P the particles belonging to the same cluster are numbered by the "internal" index $r = 1, \dots, n_\alpha$. Thus, according to Eq.(A.11),

$$\Psi_{\mathbf{q}, \mathbf{n}}^{(M)}(\mathbf{x}) = C_{\mathbf{q}, \mathbf{n}}^{(M)} \sum_P (-1)^{[P]} \left(\prod_{a < b}^N \left[-i(\partial_{x_a} - \partial_{x_b}) + i\kappa \operatorname{sgn}(x_a - x_b) \right] \right) \exp \left[i \sum_{c=1}^N q_{r(c)}^{\alpha(c)} x_c \right] \quad (\text{B.21})$$

where $C_{\mathbf{q}, \mathbf{n}}^{(M)}$ is the normalization constant to be defined later. Substituting here Eq.(B.17) and taking derivatives we get

$$\begin{aligned} \Psi_{\mathbf{q}, \mathbf{n}}^{(M)}(\mathbf{x}) &= C_{\mathbf{q}, \mathbf{n}}^{(M)} \sum_P (-1)^{[P]} \prod_{a < b}^N \left[\left(q_{\alpha(a)} - \frac{i\kappa}{2} [n_{\alpha(a)} + 1 - 2r(a)] \right) - \left(q_{\alpha(b)} - \frac{i\kappa}{2} [n_{\alpha(b)} + 1 - 2r(b)] \right) + i\kappa \operatorname{sgn}(x_a - x_b) \right] \times \\ &\times \exp \left[i \sum_{c=1}^N q_{\alpha(c)} x_c + \frac{\kappa}{2} \sum_{c=1}^N (n_{\alpha(c)} + 1 - 2r(c)) x_c \right] \end{aligned} \quad (\text{B.22})$$

The pre-exponential product in the above equation contains two types of term: the pairs of points (a, b) which belong to different clusters ($\alpha(a) \neq \alpha(b)$), and pairs of points which belong to the same cluster ($\alpha(a) = \alpha(b)$). In the last case, the product Π_α over the pairs of points which belong to a cluster Ω_α reduces to

$$\Pi_\alpha \propto \prod_{a < b \in \Omega_\alpha} [r(b) - r(a) - \operatorname{sgn}(x_a - x_b)] \quad (\text{B.23})$$

As for the ground state wave function Eq. (B.14)–(B.15), one can easily note that due to the presence of this product in the summations over $n_\alpha!$ "internal" (inside the cluster Ω_α) permutations $r(a)$ only one permutation gives non-zero contribution. To prove this statement, we note that the wave function $\Psi_{\mathbf{q}, \mathbf{n}}^{(M)}(\mathbf{x})$ is symmetric with respect to permutations of its N arguments $\{x_a\}$; it is then sufficient to consider the case where the positions of the particles are ordered, $x_1 < x_2 < \dots < x_N$. In particular, the particles $\{x_{a_k}\}$ ($k = 1, 2, \dots, n_\alpha$) belonging to the same cluster Ω_α are also ordered $x_{a_1} < x_{a_2} < \dots < x_{a_{n_\alpha}}$. In this case

$$\Pi_\alpha \propto \prod_{k < l}^{n_\alpha} [r(l) - r(k) + 1] \quad (\text{B.24})$$

Now it is evident that the above product is non-zero only for the trivial permutation, $r(k) = k$ (since if we permute any two numbers in the sequence $1, 2, \dots, n_\alpha$, we can always find two numbers $k < l$, such that $r(l) = r(k) - 1$). In this case

$$\Pi_\alpha \propto \prod_{k < l}^{n_\alpha} [l - k + 1] \quad (\text{B.25})$$

Including the values of all these "internal" products, Eq.(B.25), into the redefined normalization constant $C_{\mathbf{q},\mathbf{n}}^{(M)}$, for the wave function, Eq.(B.22) (with $x_1 < x_2 < \dots < x_N$), we obtain

$$\begin{aligned} \Psi_{\mathbf{q},\mathbf{n}}^{(M)}(\mathbf{x}) &= C_{\mathbf{q},\mathbf{n}}^{(M)} \sum_P' (-1)^{[P]} \prod_{\substack{a < b \\ \alpha(a) \neq \alpha(b)}}^N \left[\left(q_{\alpha(a)} - \frac{i\kappa}{2} n_{\alpha(a)} \right) - \left(q_{\alpha(b)} - \frac{i\kappa}{2} n_{\alpha(b)} \right) + i\kappa (r(a) - r(b) - 1) \right] \times \\ &\times \exp \left[i \sum_{c=1}^N q_{\alpha(c)} x_c + \frac{\kappa}{2} \sum_{c=1}^N (n_{\alpha(c)} + 1 - 2r(c)) x_c \right] \end{aligned} \quad (\text{B.26})$$

where the product goes now only over the pairs of particles belonging to *different* clusters, and the symbol $\sum_P'$ means that the summation goes only over the permutations P in which the "internal" indices $r(a)$ are ordered inside each cluster. Note that although the positions of particles belonging to the same cluster are ordered, the mutual positions of particles belonging to different clusters could be arbitrary, so that geometrically the clusters are free to "penetrate" each other. In other words, the name "cluster" does to assume geometrically compact particles positions.

Now taking into account the symmetry of the wave function $\Psi_{\mathbf{q},\mathbf{n}}^{(M)}(\mathbf{x})$ with respect to the permutations of its arguments the expression in Eq.(B.26) can be easily continued beyond the sector $x_1 < x_2 < \dots < x_N$ for the entire coordinate space R_N . Using the relations

$$\sum_{a \in \Omega_\alpha} (n_\alpha + 1 - 2r(a)) x_a = \sum_{k=1}^{n_\alpha} (n_\alpha + 1 - 2k) x_{a_k} = -\frac{1}{2} \sum_{k,l=1}^{n_\alpha} |x_{a_k} - x_{a_l}| \quad (\text{B.27})$$

(where $x_{a_1} < x_{a_2} < \dots < x_{a_{n_\alpha}}$), for the wave function $\Psi_{\mathbf{q},\mathbf{n}}^{(M)}(\mathbf{x})$ with *arbitrary* particles positions we get the following sufficiently compact representation (cf. Eq.(B.21)):

$$\Psi_{\mathbf{q},\mathbf{n}}^{(M)}(\mathbf{x}) = C_{\mathbf{q},\mathbf{n}}^{(M)} \sum_P' (-1)^{[P]} \prod_{\substack{a < b \\ \alpha(a) \neq \alpha(b)}}^N \left[-i(\partial_{x_a} - \partial_{x_b}) + i\kappa \operatorname{sgn}(x_a - x_b) \right] \exp \left[i \sum_{\alpha=1}^M q_\alpha \sum_{c \in \Omega_\alpha}^{n_\alpha} x_c - \frac{\kappa}{4} \sum_{\alpha=1}^M \sum_{c,c' \in \Omega_\alpha}^{n_\alpha} |x_c - x_{c'}| \right] \quad (\text{B.28})$$

Finally, substituting Eq.(B.17)-(B.18) into Eq.(A.13), for the energy spectrum one easily obtains:

$$E_M(\mathbf{q}, \mathbf{n}) = \frac{1}{2\beta} \sum_{\alpha=1}^M \sum_{r=1}^{n_\alpha} (q_\alpha^r)^2 = \frac{1}{2\beta} \sum_{\alpha=1}^M n_\alpha q_\alpha^2 - \frac{\kappa^2}{24\beta} \sum_{\alpha=1}^M (n_\alpha^3 - n_\alpha) \quad (\text{B.29})$$

3. Orthonormality

We define the overlap of two wave functions characterized by two sets of parameters, $(M, \mathbf{n}, \mathbf{q})$ and $(M', \mathbf{n}', \mathbf{q}')$ as

$$Q_{\mathbf{n},\mathbf{n}'}^{(M,M')}(\mathbf{q}, \mathbf{q}') \equiv \int_{-\infty}^{+\infty} d^N \mathbf{x} \Psi_{\mathbf{q}',\mathbf{n}'}^{(M')*}(\mathbf{x}) \Psi_{\mathbf{q},\mathbf{n}}^{(M)}(\mathbf{x}) \quad (\text{B.30})$$

Substituting here Eq.(B.28) we get

$$\begin{aligned} Q_{\mathbf{n},\mathbf{n}'}^{(M,M')}(\mathbf{q}, \mathbf{q}') &= C_{\mathbf{q},\mathbf{n}}^{(M)} C_{\mathbf{q}',\mathbf{n}'}^{(M')*} \sum_P' \sum_{P'}' (-1)^{[P]+[P']} \int_{-\infty}^{+\infty} d^N \mathbf{x} \\ &\times \left(\prod_{\substack{a < b \\ \alpha'(a) \neq \alpha'(b)}}^N \left[i(\partial_{x_a} - \partial_{x_b}) - i\kappa \operatorname{sgn}(x_a - x_b) \right] \right) \exp \left[-i \sum_{\alpha=1}^{M'} q'_\alpha \sum_{c \in \Omega'_\alpha}^{n'_\alpha} x_c - \frac{\kappa}{4} \sum_{\alpha=1}^{M'} \sum_{c,c' \in \Omega'_\alpha}^{n'_\alpha} |x_c - x_{c'}| \right] \times \\ &\times \left(\prod_{\substack{a < b \\ \alpha(a) \neq \alpha(b)}}^N \left[-i(\partial_{x_a} - \partial_{x_b}) + i\kappa \operatorname{sgn}(x_a - x_b) \right] \right) \exp \left[i \sum_{\alpha=1}^M q_\alpha \sum_{c \in \Omega_\alpha}^{n_\alpha} x_c - \frac{\kappa}{4} \sum_{\alpha=1}^M \sum_{c,c' \in \Omega_\alpha}^{n_\alpha} |x_c - x_{c'}| \right] \end{aligned} \quad (\text{B.31})$$

where $\{\Omega_\alpha\}$ and $\{\Omega'_\alpha\}$ denote the clusters of the permutations P and P' correspondingly. Integrating by parts we obtain

$$\begin{aligned}
Q_{\mathbf{n}, \mathbf{n}'}^{(M, M')}(\mathbf{q}, \mathbf{q}') &= C_{\mathbf{q}, \mathbf{n}}^{(M)} C_{\mathbf{q}', \mathbf{n}'}^{(M')*} \sum_P' \sum_{P'}' (-1)^{[P]+[P']} \int_{-\infty}^{+\infty} d^N \mathbf{x} \exp \left[-i \sum_{\alpha=1}^{M'} q'_\alpha \sum_{c \in \Omega'_\alpha}^{n'_\alpha} x_c - \frac{\kappa}{4} \sum_{\alpha=1}^{M'} \sum_{c, c' \in \Omega'_\alpha}^{n'_\alpha} |x_c - x_{c'}| \right] \times \\
&\times \left(\prod_{\substack{a < b \\ \alpha'(a) \neq \alpha'(b)}}^N \left[-i(\partial_{x_a} - \partial_{x_b}) - i\kappa \operatorname{sgn}(x_a - x_b) \right] \right) \left(\prod_{\substack{a < b \\ \alpha(a) \neq \alpha(b)}}^N \left[-i(\partial_{x_a} - \partial_{x_b}) + i\kappa \operatorname{sgn}(x_a - x_b) \right] \right) \times \\
&\times \exp \left[i \sum_{\alpha=1}^M q_\alpha \sum_{c \in \Omega_\alpha}^{n_\alpha} x_c - \frac{\kappa}{4} \sum_{\alpha=1}^M \sum_{c, c' \in \Omega_\alpha}^{n_\alpha} |x_c - x_{c'}| \right] \quad (\text{B.32})
\end{aligned}$$

First, let us consider the case when the integer parameters of the two functions coincide, $M = M'$, $\mathbf{n} = \mathbf{n}'$, and for the moment let us suppose that all these integer parameters $\{n_\alpha\}$ are *different*, $1 \leq n_1 < n_2 < \dots < n_M$. Then, in the summations over the permutations in Eq.(B.32), we find two types of terms:

- (A) the "diagonal" ones in which the two permutations coincide, $P = P'$;
- (B) the "off-diagonal" ones in which the two permutations are different, $P \neq P'$.

The contribution of the "diagonal" ones reads

$$\begin{aligned}
Q_{\mathbf{n}, \mathbf{n}}^{(M, M)^{(A)}}(\mathbf{q}, \mathbf{q}') &= C_{\mathbf{q}, \mathbf{n}}^{(M)} C_{\mathbf{q}', \mathbf{n}}^{(M)*} \sum_P' \int_{-\infty}^{+\infty} d^N \mathbf{x} \exp \left[-i \sum_{\alpha=1}^M q'_\alpha \sum_{c \in \Omega_\alpha}^{n_\alpha} x_c - \frac{\kappa}{4} \sum_{\alpha=1}^M \sum_{c, c' \in \Omega_\alpha}^{n_\alpha} |x_c - x_{c'}| \right] \times \\
&\times \left(\prod_{\substack{a < b \\ \alpha(a) \neq \alpha(b)}}^N \left[-(\partial_{x_a} - \partial_{x_b})^2 + \kappa^2 \right] \right) \exp \left[i \sum_{\alpha=1}^M q_\alpha \sum_{c \in \Omega_\alpha}^{n_\alpha} x_c - \frac{\kappa}{4} \sum_{\alpha=1}^M \sum_{c, c' \in \Omega_\alpha}^{n_\alpha} |x_c - x_{c'}| \right] \quad (\text{B.33})
\end{aligned}$$

It is evident that all permutations $\alpha(a)$ in the above equation give the same contribution and therefore it is sufficient to consider only the contribution of the "trivial" permutation which is represented by line in Eq.(B.19). The cluster ordering given by this permutation we denote by $\alpha_0(a)$. For this particular configuration of clusters we can redefine the particles numbering, so that instead of a "plane" index $a = 1, 2, \dots, N$ the particles would be counted by two indices (α, r) : $\{x_a\} \rightarrow \{x_r^\alpha\}$ ($\alpha = 1, \dots, M$) ($r = 1, \dots, n_\alpha$) indicating to which cluster α a given particle belongs and what is its "internal" cluster number r . Due to the symmetry of the integrated expression in Eq.(B.33) with respect to the permutations of the particles inside the clusters, we can introduce the "internal" particles ordering for every cluster: $x_1^\alpha < x_2^\alpha < \dots < x_{n_\alpha}^\alpha$. In this way, using the relation, Eq.(B.27), we get

$$\begin{aligned}
Q_{\mathbf{n}, \mathbf{n}}^{(M, M)^{(A)}}(\mathbf{q}, \mathbf{q}') &= C_{\mathbf{q}, \mathbf{n}}^{(M)} C_{\mathbf{q}', \mathbf{n}}^{(M)*} \frac{N!}{n_1! n_2! \dots n_M!} \left[\prod_{\alpha=1}^M \left(n_\alpha! \int_{-\infty}^{+\infty} dx_1^\alpha \int_{x_1^\alpha}^{+\infty} dx_2^\alpha \dots \int_{x_{n_\alpha-1}^\alpha}^{+\infty} dx_{n_\alpha}^\alpha \right) \right] \times \\
&\times \exp \left[-i \sum_{\alpha=1}^M q'_\alpha \sum_{r=1}^{n_\alpha} x_r^\alpha + \frac{\kappa}{2} \sum_{\alpha=1}^M \sum_{r=1}^{n_\alpha} (n_\alpha + 1 - 2r) x_r^\alpha \right] \times \\
&\times \left(\prod_{\alpha < \beta}^M \prod_{r=1}^{n_\alpha} \prod_{r'=1}^{n_\beta} \left[-(\partial_{x_r^\alpha} - \partial_{x_{r'}^\beta})^2 + \kappa^2 \right] \right) \exp \left[i \sum_{\alpha=1}^M q_\alpha \sum_{r=1}^{n_\alpha} x_r^\alpha + \frac{\kappa}{2} \sum_{\alpha=1}^M \sum_{r=1}^{n_\alpha} (n_\alpha + 1 - 2r) x_r^\alpha \right] \quad (\text{B.34})
\end{aligned}$$

where the factor $N!/n_1! \dots n_M!$ is the total number of permutations of M clusters over N particles. Taking the derivatives and reorganizing the terms we obtain

$$\begin{aligned}
Q_{\mathbf{n}, \mathbf{n}}^{(M, M)^{(A)}}(\mathbf{q}, \mathbf{q}') &= C_{\mathbf{q}, \mathbf{n}}^{(M)} C_{\mathbf{q}', \mathbf{n}}^{(M)*} N! \left(\prod_{\alpha < \beta}^M \prod_{r=1}^{n_\alpha} \prod_{r'=1}^{n_\beta} \left| \left(q_\alpha - \frac{i\kappa}{2} n_\alpha \right) - \left(q_\beta - \frac{i\kappa}{2} n_\beta \right) + i\kappa (r - r' - 1) \right|^2 \right) \times \\
&\times \prod_{\alpha=1}^M \left\{ \int_{-\infty}^{+\infty} dx_1^\alpha \int_{x_1^\alpha}^{+\infty} dx_2^\alpha \dots \int_{x_{n_\alpha-1}^\alpha}^{+\infty} dx_{n_\alpha}^\alpha e^{i(q_\alpha - q'_\alpha) \sum_{r=1}^{n_\alpha} x_r^\alpha + \kappa \sum_{r=1}^{n_\alpha} (n_\alpha + 1 - 2r) x_r^\alpha} \right\} \quad (\text{B.35})
\end{aligned}$$

Simple integrations over x_r^α yields (cf. Eqs.(B.8)-(B.10))

$$Q_{\mathbf{n},\mathbf{n}}^{(M,M)^{(A)}}(\mathbf{q},\mathbf{q}') = |C_{\mathbf{q},\mathbf{n}}^{(M)}|^2 N! \left(\prod_{\alpha < \beta}^M \prod_{r=1}^{n_\alpha} \prod_{r'=1}^{n_\beta} \left| \left(q_\alpha - \frac{i\kappa}{2} n_\alpha \right) - \left(q_\beta - \frac{i\kappa}{2} n_\beta \right) + i\kappa (r - r' - 1) \right| \right)^2 \times \\ \times \prod_{\alpha=1}^M \left[\frac{n_\alpha \kappa}{(n_\alpha!)^2 \kappa^{n_\alpha}} (2\pi) \delta(q_\alpha - q'_\alpha) \right] \quad (\text{B.36})$$

Now let us prove that the "off-diagonal" terms of Eq.(B.32), in which the permutations P and P' are different, give no contribution. Here we can also chose one of the permutations, say the permutation P , to be the "trivial" one represented by line in Eq.(B.19) with the cluster ordering denoted by $\alpha_0(a)$. Given the symmetry of the wave functions it will be sufficient to consider the contribution of the sector $x_1 < x_2 < \dots < x_N$. According to Eq.(B.32), we get

$$Q_{\mathbf{n},\mathbf{n}}^{(M,M)^{(B)}}(\mathbf{q},\mathbf{q}') \propto \sum_{P'}' (-1)^{[P']} \int_{x_1 < \dots < x_N} d^N \mathbf{x} \exp \left[-i \sum_{\alpha=1}^M q'_\alpha \sum_{a \in \Omega'_\alpha}^{n_\alpha} x_a - \frac{\kappa}{4} \sum_{\alpha=1}^M \sum_{a,b \in \Omega'_\alpha}^{n_\alpha} |x_a - x_b| \right] \times \\ \times \left(\prod_{\substack{a < b \\ \alpha_0(a) \neq \alpha_0(b)}}^N \left[-i(\partial_{x_a} - \partial_{x_b}) + i\kappa \operatorname{sgn}(x_a - x_b) \right] \right) \left(\prod_{\substack{a < b \\ \alpha'(a) \neq \alpha'(b)}}^N \left[-i(\partial_{x_a} - \partial_{x_b}) - i\kappa \operatorname{sgn}(x_a - x_b) \right] \right) \\ \times \exp \left[i \sum_{\alpha=1}^M q_\alpha \sum_{a \in \Omega_\alpha}^{n_\alpha} x_a + \frac{\kappa}{2} \sum_{\alpha=1}^M \sum_{a \in \Omega_\alpha}^{n_\alpha} (n_\alpha + 1 - 2r(a)) x_a \right] \quad (\text{B.37})$$

Here the symbols $\{\Omega_\alpha^o\}$ denote the clusters of the trivial permutation $\alpha_0(a)$. Since $P' \neq P$, some of the clusters Ω'_α must be different from Ω_α^o . As an illustration, let us consider a particular case of $N = 10$, with three clusters $n_1 = 5$ (denoted by the symbol " $\circ$ ") , $n_2 = 2$ (denoted by the symbol " $\times$ ") and $n_3 = 3$ (denoted by the symbol " $\triangle$ "):

particle number a	1	2	3	4	5	6	7	8	9	10
permutation $\alpha_0(a)$	$\circ$	$\circ$	$\circ$	$\circ$	$\circ$	$\times$	$\times$	$\triangle$	$\triangle$	$\triangle$
permutation $\alpha'(a)$	$\circ$	$\circ$	$\circ$	$\triangle$	$\circ$	$\times$	$\times$	$\circ$	$\triangle$	$\triangle$

Here in the permutation $\alpha'(a)$ the particle $a = 4$ belong to the cluster $\alpha = 3$ (and not to the cluster $\alpha = 1$ as in the permutation $\alpha_0(a)$), and the particle $a = 8$ belong to the cluster $\alpha = 1$ (and not to the cluster $\alpha = 3$ as in the permutation $\alpha_0(a)$). Now let us look carefully at the structure of the products in Eq.(B.37). Unlike the first product, which contains no "internal" products among particles belonging to the cluster Ω_1^o , the second product *does*. Besides, the signs of the differential operators $(\partial_{x_a} - \partial_{x_b})$ in the second product is *opposite* to the "normal" ones in the first product (cf. Eqs.(B.23)-(B.25)). It is these two factors (the presence of the "internal" products and the "wrong" signs of the differential operators) which makes the "off-diagonal" contributions, Eq.(B.37), to be zero. Indeed, in the above example, the second product contains the term

$$\Pi'_{4,5} \equiv \left[-i(\partial_{x_4} - \partial_{x_5}) + i\kappa \right] \exp \left[i \sum_{\alpha=1}^3 q_\alpha \sum_{a \in \Omega_\alpha^o}^{n_\alpha} x_a + \frac{\kappa}{2} \sum_{\alpha=1}^3 \sum_{a \in \Omega_\alpha^o}^{n_\alpha} (n_\alpha + 1 - 2r(a)) x_a \right] \quad (\text{B.38})$$

(we remind that the particles in the clusters Ω_α^o are ordered, and in particular $x_4 < x_5$). Taking the derivatives, we get

$$\Pi'_{4,5} \propto \left[- \left(i q_1 + \frac{\kappa}{2} (n_1 + 1 - 2r(4)) - i q_1 - \frac{\kappa}{2} (n_1 + 1 - 2r(5)) \right) + \kappa \right] \\ \propto [r(4) - r(5) + 1] = 0 \quad (\text{B.39})$$

since in the first cluster $r(a) = a$.

One can easily understand that the above example reflect the general situation. Since all the cluster sizes n_α are supposed to be different, whatever the permutation $\alpha'(a)$ is, we can always find a cluster Ω_α^o such that some of its particles belong to the same cluster number α in the permutation $\alpha'(a)$ while the others do not. Then one has to consider the contribution of the product of two *neighboring number points*

$$\Pi'_{k,k+1} = \left[-i(\partial_{x_k} - \partial_{x_{k+1}}) + i\kappa \right] \exp \left[i \sum_{\alpha=1}^M q_\alpha \sum_{a \in \Omega_\alpha^o} x_a + \frac{\kappa}{2} \sum_{\alpha=1}^M \sum_{a \in \Omega_\alpha^o} (n_\alpha + 1 - 2r(a)) x_a \right] \quad (\text{B.40})$$

where in the permutation $\alpha'(a)$ the particle k belong to the cluster number α and the particle $(k+1)$ belong to some other cluster. Taking the derivatives one gets

$$\Pi'_{k,k+1} \propto [r(k) - r(k+1) + 1] = 0 \quad (\text{B.41})$$

as $r(a)$ is the "internal" particle number in the cluster Ω_α^o , where $r(k+1) = r(k) + 1$ (cf. Eqs.(B.23)-(B.25)).

Thus, the only non-zero contribution to the overlap, Eq.(B.30), of two wave function $\Psi_{\mathbf{q}', \mathbf{n}}^{(M)}(\mathbf{x})$ and $\Psi_{\mathbf{q}, \mathbf{n}}^{(M)}(\mathbf{x})$ (having the same number of clusters M and characterized by the same set of the integer parameters $1 \leq n_1 < n_2 < \dots < n_M$) comes from the "diagonal" terms, Eq.(B.36):

$$\begin{aligned} Q_{\mathbf{n}, \mathbf{n}}^{(M, M)}(\mathbf{q}, \mathbf{q}') &= |C_{\mathbf{q}, \mathbf{n}}^{(M)}|^2 N! \prod_{\alpha=1}^M \left[\frac{n_\alpha \kappa}{(n_\alpha!)^2 \kappa^{n_\alpha}} \right] \left(\prod_{\alpha < \beta} \prod_{r=1}^{n_\alpha} \prod_{r'=1}^{n_\beta} \left| \left(q_\alpha - \frac{i\kappa}{2} n_\alpha \right) - \left(q_\beta - \frac{i\kappa}{2} n_\beta \right) + i\kappa (r - r' - 1) \right|^2 \right) \times \\ &\times \prod_{\alpha=1}^M \left[(2\pi) \delta(q_\alpha - q'_\alpha) \right] \end{aligned} \quad (\text{B.42})$$

The situation when there are clusters which have the same numbers of particles n_α is somewhat more complicated. Let us consider the overlap between two wave function $\Psi_{\mathbf{q}', \mathbf{n}}^{(M)}(\mathbf{x})$ and $\Psi_{\mathbf{q}, \mathbf{n}}^{(M)}(\mathbf{x})$ (which, as before have the same M and $\mathbf{n}$) such that in the set of M integers $n_1, n_2, \dots, n_M$ there are two n_α 's which are equal, say $n_{\alpha_1} = n_{\alpha_2}$ (where $\alpha_1 \neq \alpha_2$). In the eigenstate $(\mathbf{q}', \mathbf{n})$ these two clusters have the center of mass momenta q'_{α_1} and q'_{α_2} , and in the the eigenstate $(\mathbf{q}, \mathbf{n})$ they have the momenta q_{α_1} and q_{α_2} correspondingly. According to the above discussion, the non-zero contributions in the summation over the cluster permutations $\alpha(a)$ and $\alpha'(a)$ in Eq.(B.32) appears only if the clusters $\{\Omega_\alpha\}$ of the permutation $\alpha(a)$ totally coincide with the clusters $\{\Omega'_\alpha\}$ of the permutation $\alpha'(a)$. In the case when all n_α are different this is possible only if the permutation $\alpha(a)$ coincides with the permutation $\alpha'(a)$. In contrast to that, in the case when we have $n_{\alpha_1} = n_{\alpha_2}$, there are *two* non-zero options. The first one, as before, is given by the "diagonal" terms with $\alpha(a) = \alpha'(a)$ (so that the clusters $\{\Omega_\alpha\}$ and $\{\Omega'_\alpha\}$ are just the same), and this contribution is proportional to $\delta(q_{\alpha_1} - q'_{\alpha_1}) \delta(q_{\alpha_2} - q'_{\alpha_2})$. The second ("off-diagonal") contribution is given by such permutation $\alpha'(a)$ in which the cluster Ω'_{α_1} (of the permutation $\alpha'(a)$) coincide with the cluster Ω_{α_2} (of the permutation $\alpha(a)$) and the cluster Ω'_{α_2} (of the permutation $\alpha'(a)$) coincide with the cluster Ω_{α_1} (of the permutation $\alpha(a)$) while the rest of the clusters of these two permutations are the same, $\Omega'_\alpha = \Omega_\alpha$ ($\alpha \neq \alpha_1, \alpha_2$). Correspondingly, this last contribution is proportional to $\delta(q_{\alpha_1} - q'_{\alpha_2}) \delta(q_{\alpha_2} - q'_{\alpha_1}) (-1)^{n_{\alpha_1}}$. In fact this situation with two equivalent contributions is the consequence of the symmetry of the wave function $\Psi_{\mathbf{q}, \mathbf{n}}^{(M)}(\mathbf{x})$: the permutation of two momenta q_{α_1} and q_{α_2} belonging to the clusters which have the same numbers of particles, $n_{\alpha_1} = n_{\alpha_2}$ produces the factor $(-1)^{n_{\alpha_1}}$. This is evident from the general expression for the wave function, eq.(A.9), where the permutation of any two momenta q_{α_1} and q_{α_2} belonging to the clusters which have the same numbers of particles corresponds to the permutation of n columns of the matrix $\exp(iq_a x_b)$. Therefore considering the clusters with equal numbers of particles as equivalent and restricting analysis to the sectors $q_{\alpha_1} < q_{\alpha_2}$; $q'_{\alpha_1} < q'_{\alpha_2}$ we find that the second contribution, $\delta(q_{\alpha_1} - q'_{\alpha_2}) \delta(q_{\alpha_2} - q'_{\alpha_1})$ is identically equal to zero, thus returning to the above result Eq.(B.42).

A generic eigenstate $(\mathbf{q}, \mathbf{n})$ with M clusters could be specified in terms of the following set of parameters:

$$(\mathbf{q}, \mathbf{n}) = \{ \underbrace{(q_1, m_1), \dots, (q_{s_1}, m_1)}_{s_1}, \underbrace{(q_{s_1+1}, m_2), \dots, (q_{s_1+s_2}, m_2)}_{s_2}, \dots; \underbrace{(q_{s_1+\dots+s_{k-1}+1}, m_k), \dots, (q_{s_1+\dots+s_k}, m_k)}_{s_k} \} \quad (\text{B.43})$$

where $s_1 + s_2 + \dots + s_k = M$ and k integers $\{m_i\}$ ($1 \leq k \leq M$) are all supposed to be *different*:

$$1 \leq m_1 < m_2 < \dots < m_k \quad (\text{B.44})$$

Here the integer parameter k denotes the number of different cluster types. For a given k

$$\sum_{\alpha=1}^M n_\alpha = \sum_{i=1}^k s_i m_i = N \quad (\text{B.45})$$

Due to the symmetry with respect to the momenta permutations inside the subsets of equal n 's it is sufficient to consider the wave functions in the sectors

$$\begin{aligned} q_1 &< q_2 < \dots < q_{s_1} ; \\ q_{s_1+1} &< q_{s_1+2} < \dots < q_{s_1+s_2} ; \\ &\dots\dots\dots \\ q_{s_1+\dots+s_{k-1}+1} &< q_{s_1+\dots+s_{k-1}+2} < \dots < q_{s_1+\dots+s_{k-1}+s_k} \end{aligned} \quad (\text{B.46})$$

In this representation we again recover the above result Eq.(B.42)

Finally, let us consider the overlap of two eigenstates described by two *different* sets of integer parameters, $\mathbf{n}' \neq \mathbf{n}$. In fact this situation is quite simple because if the clusters of the two states are different from each other, it means that in the summation over the pairs of permutations P and P' in Eq.(B.32) there exist no two permutations for which these two sets of clusters $\{\Omega_\alpha\}$ and $\{\Omega'_\alpha\}$ would coincide. Which, according to the above analysis, means that this expression is equal to zero. Note that the condition $M' \neq M$ automatically implies that $\mathbf{n}' \neq \mathbf{n}$.

Thus we have proved that

$$\begin{aligned} Q_{\mathbf{n}, \mathbf{n}'}^{(M, M')}(\mathbf{q}, \mathbf{q}') &= |C_{\mathbf{q}, \mathbf{n}}^{(M)}|^2 \delta(M, M') \left(\prod_{\alpha=1}^M \delta(n_\alpha, n'_\alpha) \right) \left(\prod_{\alpha=1}^M (2\pi) \delta(q_\alpha - q'_\alpha) \right) \times \\ &\times N! \prod_{\alpha=1}^M \left[\frac{n_\alpha \kappa}{(n_\alpha!)^2 \kappa^{n_\alpha}} \right] \prod_{\alpha < \beta}^M \prod_{r=1}^{n_\alpha} \prod_{r'=1}^{n_\beta} \left| \left(q_\alpha - \frac{i\kappa}{2} n_\alpha \right) - \left(q_\beta - \frac{i\kappa}{2} n_\beta \right) + i\kappa (r - r' - 1) \right|^2 \end{aligned} \quad (\text{B.47})$$

where the integer parameters $\{n_\alpha\}$ and $\{n'_\alpha\}$ are assumed to have the generic structure represented in Eqs.(B.43)-(B.45), and the momenta $\{q_\alpha\}$ and $\{q'_\alpha\}$ of the clusters with equal numbers of particles are restricted in the sectors, Eq.(B.46). According to Eq.(B.47), the orthonormality condition defines the normalization constant

$$|C^{(M)}(\mathbf{q}, \mathbf{n})|^2 = \frac{1}{N!} \left[\prod_{\alpha=1}^M \frac{(n_\alpha!)^2 \kappa^{n_\alpha}}{n_\alpha \kappa} \right] \prod_{\alpha < \beta}^M \prod_{r=1}^{n_\alpha} \prod_{r'=1}^{n_\beta} \frac{1}{\left| \left(q_\alpha - \frac{i\kappa}{2} n_\alpha \right) - \left(q_\beta - \frac{i\kappa}{2} n_\beta \right) + i\kappa (r - r' - 1) \right|^2} \quad (\text{B.48})$$

In other words, the wave functions, Eq.(B.28), form the orthonormal set. Although, at present we are not able to prove that this set is complete, the *suggestion* of the completeness (which assumes that there are exist no other eigenstates besides those described above) looks quite natural.

4. Propagator

The time dependent solution $\Psi(\mathbf{x}, t)$ of the imaginary-time Schrödinger equation

$$\beta \partial_t \Psi(\mathbf{x}; t) = \frac{1}{2} \sum_{a=1}^N \partial_{x_a}^2 \Psi(\mathbf{x}; t) + \frac{1}{2} \kappa \sum_{a \neq b}^N \delta(x_a - x_b) \Psi(\mathbf{x}; t) \quad (\text{B.49})$$

with the initial condition

$$\Psi(\mathbf{x}; 0) = \prod_{a=1}^N \delta(x_a) \quad (\text{B.50})$$

can be represented in terms of the linear combination of the eigenfunctions $\Psi_{\mathbf{q}, \mathbf{n}}^{(M)}(\mathbf{x})$, Eq.(B.28):

$$\Psi(\mathbf{x}, t) = \sum_{M=1}^N \sum'_{\mathbf{n}} \int' \mathcal{D}\mathbf{q} \Psi_{\mathbf{q}, \mathbf{n}}^{(M)}(\mathbf{x}) \Psi_{\mathbf{q}, \mathbf{n}}^{(M)*}(\mathbf{0}) \exp[-E_M(\mathbf{q}, \mathbf{n}) t] \quad (\text{B.51})$$

where the energy spectrum $E_M(\mathbf{q}, \mathbf{n})$ is given by Eq.(B.29). The summations over n_α are performed here in terms of the parameters $\{s_i, m_i\}$, Eqs.(B.43)-(B.45):

$$\sum'_{\mathbf{n}} \equiv \sum_{k=1}^M \sum_{s_1 \dots s_k=1}^{\infty} \sum_{1 \leq m_1 \dots < m_k}^{\infty} \delta\left(\sum_{i=1}^k s_i, M\right) \delta\left(\sum_{i=1}^k s_i m_i, N\right) \quad (\text{B.52})$$

where $\delta(n, l)$ is the Kronecker symbol, and for simplicity (due to the presence of these Kronecker symbols) the summations over m_i and s_i are extended to infinity. The symbol $\int' \mathcal{D}\mathbf{q}$ in Eq.(B.51) denotes the integration over M momenta q_α in the sectors, Eq.(B.46).

The replica partition function $Z(N, L)$ of the original directed polymer problem is obtained via a particular choice of the final-point coordinates,

$$Z(N, L) = \Psi(\mathbf{0}; L) = \sum_{M=1}^N \sum'_{\mathbf{n}} \int' \mathcal{D}\mathbf{q} \left| \Psi_{\mathbf{q}, \mathbf{n}}^{(M)}(\mathbf{0}) \right|^2 \exp[-E_M(\mathbf{q}, \mathbf{n}) L] \quad (\text{B.53})$$

According to Eq.(B.28), for $M \geq 2$,

$$\begin{aligned} \Psi_{\mathbf{q}, \mathbf{n}}^{(M)}(\mathbf{0}) &= C_{\mathbf{q}, \mathbf{n}}^{(M)} \sum'_P (-1)^{[P]} \prod_{\substack{a < b \\ \alpha(a) \neq \alpha(b)}}^N \left[\left(q_{\alpha(a)} - \frac{i\kappa}{2} [n_{\alpha(a)} + 1 - 2r(a)] \right) - \left(q_{\alpha(b)} - \frac{i\kappa}{2} [n_{\alpha(b)} + 1 - 2r(b)] \right) \right] \\ &= C_{\mathbf{q}, \mathbf{n}}^{(M)} \frac{N!}{n_1! n_2! \dots n_M!} \prod_{\alpha < \beta}^M \prod_{r=1}^{n_\alpha} \prod_{r'=1}^{n_\beta} \left[\left(q_\alpha - \frac{i\kappa}{2} n_\alpha \right) - \left(q_\beta - \frac{i\kappa}{2} n_\beta \right) + i\kappa (r - r') \right] \end{aligned} \quad (\text{B.54})$$

Substituting here the value of the normalization constant, Eq.(B.48), we get

$$\left| \Psi_{\mathbf{q}, \mathbf{n}}^{(M)}(\mathbf{0}) \right|^2 = \frac{N! \kappa^N}{\left(\prod_{\alpha=1}^M \kappa n_\alpha \right)} \prod_{\alpha < \beta}^M \frac{\prod_{r=1}^{n_\alpha} \prod_{r'=1}^{n_\beta} \left| \left(q_\alpha - \frac{i\kappa}{2} n_\alpha \right) - \left(q_\beta - \frac{i\kappa}{2} n_\beta \right) + i\kappa (r - r') \right|^2}{\prod_{r=1}^{n_\alpha} \prod_{r'=1}^{n_\beta} \left| \left(q_\alpha - \frac{i\kappa}{2} n_\alpha \right) - \left(q_\beta - \frac{i\kappa}{2} n_\beta \right) + i\kappa (r - r' - 1) \right|^2} \quad (\text{B.55})$$

This expression can be essentially simplified. Shifting the product over r' in the denominator by 1 we obtain

$$\left| \Psi_{\mathbf{q}, \mathbf{n}}^{(M)}(\mathbf{0}) \right|^2 = \frac{N! \kappa^N}{\left(\prod_{\alpha=1}^M \kappa n_\alpha \right)} \prod_{\alpha < \beta}^M \frac{\prod_{r=1}^{n_\alpha} \left| \left(q_\alpha - \frac{i\kappa}{2} n_\alpha \right) - \left(q_\beta - \frac{i\kappa}{2} n_\beta \right) + i\kappa (r - 1) \right|^2}{\prod_{r=1}^{n_\alpha} \left| \left(q_\alpha - \frac{i\kappa}{2} n_\alpha \right) - \left(q_\beta - \frac{i\kappa}{2} n_\beta \right) + i\kappa (r - n_\beta - 1) \right|^2} \quad (\text{B.56})$$

Redefining the product parameter r in the denominator, $r \rightarrow n_\alpha + 1 - r$, and changing the obtained expression (under the modulus square) by its complex conjugate we get

$$\left| \Psi_{\mathbf{q}, \mathbf{n}}^{(M)}(\mathbf{0}) \right|^2 = \frac{N! \kappa^N}{\left(\prod_{\alpha=1}^M \kappa n_\alpha \right)} \prod_{\alpha < \beta}^M \frac{\prod_{r=1}^{n_\alpha} \left| \left(q_\alpha - \frac{i\kappa}{2} n_\alpha \right) - \left(q_\beta - \frac{i\kappa}{2} n_\beta \right) + i\kappa (r - 1) \right|^2}{\prod_{r=1}^{n_\alpha} \left| \left(q_\alpha - \frac{i\kappa}{2} n_\alpha \right) - \left(q_\beta - \frac{i\kappa}{2} n_\beta \right) + i\kappa r \right|^2} \quad (\text{B.57})$$

Shifting now the product over r in the numerator by 1 we finally obtain

$$\left| \Psi_{\mathbf{q}, \mathbf{n}}^{(M)}(\mathbf{0}) \right|^2 = \frac{N! \kappa^N}{\left(\prod_{\alpha=1}^M \kappa n_\alpha \right)} \prod_{\alpha < \beta}^M \frac{\left| q_\alpha - q_\beta - \frac{i\kappa}{2} (n_\alpha - n_\beta) \right|^2}{\left| q_\alpha - q_\beta - \frac{i\kappa}{2} (n_\alpha + n_\beta) \right|^2} \quad (\text{B.58})$$

For $M = 1$, according to Eqs.(B.1) and (B.11),

$$\left| \Psi_q^{(1)}(\mathbf{0}) \right|^2 = \frac{\kappa^N N!}{\kappa N} \quad (\text{B.59})$$

Since the function $f(\mathbf{q}, \mathbf{n}) = \left| \Psi_{\mathbf{q}, \mathbf{n}}^{(M)}(\mathbf{0}) \right|^2 \exp[-E_M(\mathbf{q}, \mathbf{n}) L]$ in Eq.(B.53) is symmetric with respect to permutations of all its M pairs of arguments (q_α, n_α) the integrations over M momenta q_α can be extended beyond the sector defined in Eq.(B.46) for the whole space R_M . As a consequence, there is no need to distinguish equal and different n_α 's any more, and instead of Eq.(B.52), we can sum over M integer parameters n_α with the only constrain, Eq.(B.18) (note that this kind of simplifications holds only for the specific "zero final-point" object $\Psi(\mathbf{0}; t)$, Eq.(B.53), and *not* for the general propagator $\Psi(\mathbf{x}; t)$, Eq.(B.51) containing N arbitrary coordinates $x_1, \dots, x_N$). Thus, instead of Eq.(B.53) we get

$$Z(N, L) = \sum_{M=1}^N \frac{1}{M!} \left[\prod_{\alpha=1}^M \int_{-\infty}^{+\infty} \frac{dq_\alpha}{2\pi} \sum_{n_\alpha=1}^{\infty} \right] \delta\left(\sum_{\alpha=1}^M n_\alpha, N\right) \left| \Psi_{\mathbf{q}, \mathbf{n}}^{(M)}(\mathbf{0}) \right|^2 e^{-E_M(\mathbf{q}, \mathbf{n}) L} \quad (\text{B.60})$$

Substituting here Eqs.(B.29), (B.58) and (B.59) we get the following sufficiently compact representation for the replica partition function:

$$\begin{aligned}
Z(N, L) = & N! \kappa^N \left\{ \int_{-\infty}^{+\infty} \frac{dq}{2\pi\kappa N} \exp \left[-\frac{L}{2\beta} N q^2 + \frac{\kappa^2 L}{24\beta} (N^3 - N) \right] + \right. \\
& + \sum_{M=2}^N \frac{1}{M!} \left[\prod_{\alpha=1}^M \sum_{n_\alpha=1}^{\infty} \int_{-\infty}^{+\infty} \frac{dq_\alpha}{2\pi\kappa n_\alpha} \right] \delta \left(\sum_{\alpha=1}^M n_\alpha, N \right) \prod_{\alpha < \beta}^M \frac{|q_\alpha - q_\beta - \frac{i\kappa}{2}(n_\alpha - n_\beta)|^2}{|q_\alpha - q_\beta - \frac{i\kappa}{2}(n_\alpha + n_\beta)|^2} \times \\
& \times \exp \left[-\frac{L}{2\beta} \sum_{\alpha=1}^M n_\alpha q_\alpha^2 + \frac{\kappa^2 L}{24\beta} \sum_{\alpha=1}^M (n_\alpha^3 - n_\alpha) \right] \Big\}
\end{aligned} \tag{B.61}$$

The first term in the above expression is the contribution of the ground state ($M = 1$), while the next terms ($M \geq 2$) are the contributions of the rest of the energy spectrum. After simple algebra the above replica partition function can be represented as follows:

$$Z(N, L) = e^{-\beta N L f_0} \tilde{Z}(N, \lambda) \tag{B.62}$$

where $f_0 = \frac{1}{24}\beta^4 u^2 - \frac{1}{\beta L} \ln(\beta^3 u)$, and

$$\begin{aligned}
\tilde{Z}(N, L) = & N! \left\{ \int_{-\infty}^{+\infty} \frac{dq}{2\pi\kappa N} \exp \left[-\frac{L}{2\beta} N q^2 + \frac{\kappa^2 L}{24\beta} N^3 \right] + \right. \\
& + \sum_{M=2}^N \frac{1}{M!} \left[\prod_{\alpha=1}^M \sum_{n_\alpha=1}^{\infty} \int_{-\infty}^{+\infty} \frac{dq_\alpha}{2\pi\kappa n_\alpha} \right] \delta \left(\sum_{\alpha=1}^M n_\alpha, N \right) \prod_{\alpha < \beta}^M \frac{|q_\alpha - q_\beta - \frac{i\kappa}{2}(n_\alpha - n_\beta)|^2}{|q_\alpha - q_\beta - \frac{i\kappa}{2}(n_\alpha + n_\beta)|^2} \times \\
& \times \exp \left[-\frac{L}{2\beta} \sum_{\alpha=1}^M n_\alpha q_\alpha^2 + \frac{\kappa^2 L}{24\beta} \sum_{\alpha=1}^M n_\alpha^3 \right] \Big\}
\end{aligned} \tag{B.63}$$

Appendix C

Fredholm determinant with the Airy kernel and the Tracy-Widom distribution

In this Appendix the original derivation of Tracy and Widom [14] will be repeated in simple terms to demonstrate that the function $F_2(s)$ defined as the Fredholm determinant with the Airy kernel can be expressed in terms of the solution of the Panlevé II differential equation, namely

$$F_2(s) \equiv \det[1 - \hat{K}_A] = \exp \left[- \int_s^\infty dt (t-s) q^2(t) \right] \tag{C.1}$$

where $\hat{K}_A$ is the integral operator defined on semi-infinite interval $[s, \infty)$ with the Airy kernel,

$$K_A(t_1, t_2) = \frac{\text{Ai}(t_1) \text{Ai}'(t_2) - \text{Ai}'(t_1) \text{Ai}(t_2)}{t_1 - t_2} \tag{C.2}$$

and the function $q(t)$ is the solution of the Panlevé II equation,

$$q'' = tq + 2q^3 \tag{C.3}$$

with the boundary condition, $q(t \rightarrow +\infty) \sim \text{Ai}(t)$.

Let us introduce a new function $R(t)$ such that

$$F_2(s) = \exp \left[- \int_s^\infty dt R(t) \right] \tag{C.4}$$

or, according to the definition, Eq.(C.1),

$$R(s) = \frac{d}{ds} \ln \left[\det(1 - \hat{K}_A) \right] \quad (\text{C.5})$$

Here the logarithm of the determinant can be expressed in terms of the trace:

$$\begin{aligned} \ln \left[\det(1 - \hat{K}_A) \right] &= - \sum_{n=1}^{\infty} \frac{1}{n} \text{Tr} \hat{K}_A^n \\ &\equiv - \sum_{n=1}^{\infty} \frac{1}{n} \int_s^{\infty} dt_1 \int_s^{\infty} dt_2 \dots \int_s^{\infty} dt_n K_A(t_1, t_2) K_A(t_2, t_3) \dots K_A(t_n, t_1) \end{aligned} \quad (\text{C.6})$$

Taking derivative of this expression we gets

$$\begin{aligned} R(s) &= - \int_s^{\infty} dt (1 - \hat{K}_A)^{-1}(s, t) K_A(t, s) \\ &\equiv -K_A(s, s) - \sum_{n=2}^{\infty} \int_s^{\infty} dt_1 \int_s^{\infty} dt_2 \dots \int_s^{\infty} dt_{n-1} K_A(s, t_1) K_A(t_1, t_2) \dots K_A(t_{n-1}, s) \end{aligned} \quad (\text{C.7})$$

Substituting here the integral representation of the Airy kernel, Eq.(C.2),

$$K_A(t_1, t_2) = \int_0^{\infty} dz \text{Ai}(t_1 + z) \text{Ai}(t_2 + z) \quad (\text{C.8})$$

after some efforts in simple algebra one gets

$$R(s) = \int_s^{\infty} dt_1 \int_s^{\infty} dt_2 \text{Ai}(t_1) (1 - \hat{K}_A)^{-1}(t_1, t_2) \text{Ai}(t_2) \quad (\text{C.9})$$

Taking the derivative of this expression and applying some more efforts in slightly more complicated algebra, we obtain

$$\frac{d}{ds} R(s) = -q^2(s) \quad (\text{C.10})$$

where

$$q(s) = \int_s^{\infty} dt (1 - \hat{K}_A)^{-1}(s, t) \text{Ai}(t) \quad (\text{C.11})$$

According to Eq.(C.10),

$$R(s) = \int_s^{\infty} dt q^2(t) \quad (\text{C.12})$$

Let us introduce two more functions

$$v(s) = \int_s^{\infty} dt_1 \int_s^{\infty} dt_2 \text{Ai}(t_1) (1 - \hat{K}_A)^{-1}(t_1, t_2) \text{Ai}'(t_2) \quad (\text{C.13})$$

$$p(s) = \int_s^{\infty} dt (1 - \hat{K}_A)^{-1}(s, t) \text{Ai}'(t) \quad (\text{C.14})$$

Taking derivatives of the above three functions $q(s)$, $v(s)$ and $p(s)$, Eqs.(C.11), (C.13) and (C.14), after somewhat painfull algebra one finds the following three relations:

$$q' = p - Rq \quad (\text{C.15})$$

$$p' = sq - pR - 2qv \quad (\text{C.16})$$

$$v' = -pq \quad (\text{C.17})$$

Taking derivative of the combination $(R^2 - 2v)$ and using Eqs.(C.10) and (C.17), we get

$$\frac{d}{ds}(R^2 - 2v) = 2q(p - Rq) \quad (\text{C.18})$$

On the other hand, multiplying Eq.(C.15) by $2q$ we find

$$\frac{d}{ds}q^2 = 2q(p - Rq) \quad (\text{C.19})$$

Comparing Eqs.(C.18) and (C.19) and taking into account that the value of all the above functions at $s \rightarrow \infty$ is zero, we obtain the following relation

$$R^2 - 2v = q^2 \quad (\text{C.20})$$

Finally, taking the derivative of Eq.(C.15) and using Eqs.(C.10), (C.15), (C.16) and (C.20) we easily find

$$q'' = 2q^3 + sq \quad (\text{C.21})$$

which is the special case of the Panlevé II differential equation [36, 37]. Thus, substituting Eq.(C.12) into Eq.(C.4) we obtain Eq.(C.1).

The function $F_2(s)$ gives the probability that a random quantity t described by a probability distribution functions $P_{TW}(t)$ has the value bigger than a given parameter s :

$$F_2(s) = \int_s^\infty dt P_{TW}(t) \quad (\text{C.22})$$

Taking the derivative of this relation and substituting here the result, Eq.(C.1), we find

$$P_{TW}(s) = \exp\left[-\int_s^\infty dt (t-s)q^2(t)\right] \times \int_s^\infty dt q^2(t) \quad (\text{C.23})$$

In the limit $s \rightarrow \infty$ the function $q(s)$, according to its definition, eq.(C.11), must go to zero, and in this case Eq.(C.21) turns into the Airy function equation, $q'' = sq$. Thus

$$q(s \rightarrow \infty) \simeq \text{Ai}(s) \sim \exp\left[-\frac{2}{3}s^{3/2}\right] \quad (\text{C.24})$$

It can be proved [38] that in the opposite limit, $s \rightarrow -\infty$, the asymptotic form of the solution of the Panleveé equation (C.21) (which has the right tail Airy function limit, Eq.(C.24)) is

$$q(s \rightarrow -\infty) \simeq \sqrt{-\frac{1}{2}s} \quad (\text{C.25})$$

Substituting the above two asymptotics into Eq.(C.23), we can estimate the asymptotic behaviour for the right and the left tails of the TW probability distribution function:

$$P_{TW}(s \rightarrow +\infty) \sim \exp\left[-\frac{4}{3}s^{3/2}\right] \quad (\text{C.26})$$

$$P_{TW}(s \rightarrow -\infty) \sim \exp\left[-\frac{1}{12}|s|^3\right] \quad (\text{C.27})$$

-
- [1] T. Halpin-Healy and Y-C. Zhang, Phys. Rep. **254**, 215 (1995).
 - [2] S. Lemerle, J. Ferré, C. Chappert, V. Mathet, T. Giamarchi, and P. Le Doussal, Phys. Rev. Lett. **80**, 849 (1998).
 - [3] G. Blatter, M.V. Feigel'man, V.B. Geshkenbein, A.I. Larkin, and V.M. Vinokur, Rev. Mod. Phys. **66**, 1125 (1994).
 - [4] D. Wilkinson and J.F. Willemsen, J. Phys. A **16**, 3365 (1983).
 - [5] J.M. Burgers, *The Nonlinear Diffusion Equation* (Reidel, Dordrecht, 1974).
 - [6] D.A. Huse, C.L. Henley, and D.S. Fisher, Phys. Rev. Lett. **55**, 2924 (1985).
 - [7] D.A. Huse and C.L. Henley, Phys. Rev. Lett. **54**, 2708 (1985).
 - [8] M. Kardar and Y-C. Zhang, Phys. Rev. Lett. **58**, 2087 (1987).
 - [9] M. Kardar, Nucl. Phys. **B 290**, 582 (1987).
 - [10] E. Medina and M. Kardar, J. Stat. Phys. **71**, 967 (1993).
 - [11] V.S. Dotsenko, L.B. Ioffe, V.B. Geshkenbein, S.E. Korshunov and G. Blatter, Phys. Rev. Lett. **100**, 050601 (2008)
 - [12] Yi-Cheng Zhang, Europhys. Lett. **9**, 113 (1989)
 - [13] I.V. Kolokolov and S.E. Korshunov, Phys. Rev. B **75**, 140201(R) (2007); Phys. Rev. B **78**, 024206 (2008); Phys. Rev. E **80**, 031107 (2009)
 - [14] C.A. Tracy and H. Widom, Commun. Math. Phys. **159**, 151 (1994)
 - [15] M. Prähofer and H. Spohn, Phys. Rev. Lett. **84**, 4882 (2000).
 - [16] J. Baik, P.A. Deift and K. Johansson, J. Amer. Math. Soc. **12**, 1119 (1999)
 - [17] S.N. Majumdar and S. Nechaev, Phys. Rev. E **72**, 020901(R) (2005)
 - [18] J. Gravner, C.A. Tracy and H. Widom, J. Stat. Phys. **102**, 1085 (2001)
 - [19] S.N. Majumdar and S. Nechaev, Phys. Rev. E **69**, 011103 (2004)
 - [20] K. Johansson, Comm. Math. Phys. **209**, 437 (2000)
 - [21] E. Brunet and B. Derrida, Phys. Rev. E **61**, 6789 (2000).
 - [22] M. Prähofer and H. Spohn, J. Stat. Phys. **115**, 255 (2004)
 - [23] T.Sasamoto and H.Spohn, arXiv:1002.1873; arXiv:1002.1879; arXiv:1002.1883
 - [24] G.Amir, I.Corwin and J.Quastel, arXiv:1003.0443
 - [25] V.Dotsenko and B.Klumov, J.Stat.Mech. P03022 (2010)
 - [26] P.Calabrese, P. Le Doussal and A.Rosso arXiv:1002.4560
 - [27] V.Dotsenko, arXiv:1003.4899
 - [28] E.H. Lieb and W. Liniger, Phys. Rev. **130**, 1605 (1963)
 - [29] V.E. Korepin, N.M. Bogoliubov, and A.G. Izergin, *Quantum inverse scattering method and correlation functions* (Cambridge Univ. Press, Cambridge, 1993)
 - [30] M. Gaudin, *La fonction d'onde de Bethe*, (Paris, Masson, 1983)
 - [31] J.B. McGuire, J. Math. Phys. **5**, 622 (1964).
 - [32] C.N. Yang, Phys. Rev. **168**, 1920 (1968)
 - [33] M. Takahashi, *Thermodynamics of one-dimensional solvable models* (Cambridge University Press, 1999).
 - [34] P. Calabrese and J.-S. Caux, Phys. Rev. Lett. **98**, 150403 (2007).
 - [35] M.L.Mehta, *Random Matrices* (Elsevier, Amsterdam 2004)
 - [36] P.Panleveé, *Sur les équation différentielles du second ordre et ordre supérieur dont intégrale générale est uniforme*. Acta. Math. **25**, 1-85 (1902)
 - [37] K.Iwasaki, H.Kimura, S.Shimomura and M.Yoshida, *From Gauss to Panleveé: a modern theory of special functions*. (Braunschweig, Vieweg 1991)
 - [38] S.P.Hastings and J.B.McLeod, Arch. Rat. Mech. Anal. **73**, (1980)